

TAMATE

Topiramate tablets

1 NAME OF THE MEDICINE

Topiramate

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 25 mg, 50 mg, 100 mg, or 200 mg of Topiramate as the active ingredient.

For the full list of excipients, see Section 6.1 List of excipients.

3 PHARMACEUTICAL FORM

TAMATE 25 mg: White, round, biconvex, film-coated tablets debossed with “G” on one side and “TO” over “25” on the other.

TAMATE 50 mg: Yellow, round, biconvex, film-coated tablets debossed with “G” on one side and “TO” over “50” on the other.

TAMATE 100 mg: Yellow, round, biconvex, film-coated tablets debossed with “G” on one side and “TO” over “100” on the other.

TAMATE 200 mg: Red, round, biconvex, film-coated tablets debossed with “G” on one side and “TO” over “200” on the other.

4 CLINICAL PARTICULARS

4.1 THERAPEUTIC INDICATIONS

Epilepsy

TAMATE is indicated in adults and children, 2 years and over:

- as monotherapy in patients with newly diagnosed epilepsy
- for conversion to monotherapy in patients with epilepsy
- as add-on therapy in partial onset seizures (with or without secondary generalised seizures), primary generalised tonic-clonic seizures or drop attacks associated with Lennox-Gastaut syndrome.

Migraine

TAMATE is indicated for the prophylaxis of migraine headache in adults. The usefulness of topiramate in the acute treatment of migraine headache has not been studied.

4.2 DOSE AND METHOD OF ADMINISTRATION

Method of administration

TAMATE tablets should be swallowed whole.

TAMATE can be taken without regard to meals.

Dosage (dose and interval)

For optimum seizure control in both adults and children, it is recommended that therapy should be initiated at a low dose followed by slow titration to an effective dose. Dose titration should be guided by clinical outcome.

The recommended dosages of TAMATE in adults and children for epilepsy are summarised in **Table 1**.

Epilepsy - Monotherapy

In newly diagnosed epileptic patients, TAMATE monotherapy should be initiated at a low dose (see **Table 1**).

In patients who are being converted to TAMATE monotherapy, consideration should be given to the effects of seizure control when withdrawing concomitant antiepileptic agents (AEDs). Unless safety concerns require an abrupt withdrawal of the concomitant AED, a gradual discontinuation at the rate of approximately one-third of the concomitant AED dose every 2 weeks is recommended (see **Drug withdrawal and Dosage reduction** and **section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE**). When enzyme inducing drugs are withdrawn, topiramate levels will increase. A decrease in TAMATE dosage may be required if clinically indicated.

Adult

Titration for monotherapy should begin at 25 mg as a single (nightly) dose for one week or longer. The dosage should then be increased by 25 to 50 mg/day at weekly or longer intervals to the recommended target dose of 100 mg/day. If the patient is unable to tolerate the titration regimen, smaller increments or longer intervals between increments can be used. The maximum recommended dose is 500 mg/day. Some patients with refractory forms of epilepsy have tolerated doses of 1,000 mg/day. The daily dosage should be taken as two divided doses.

Children (2 years and over)

Titration for monotherapy should begin at 0.5 to 1 mg/kg as a single (nightly) dose for the first week. The dosage should then be increased by 0.5 to 1 mg/kg/day at weekly or longer intervals to the recommended target dose of 3 to 6 mg/kg/day. If the child is unable to tolerate the titration regimen, smaller increments or longer intervals between dose increments can be used. Some children with recently diagnosed partial onset seizures have received doses of up to 500 mg/day. The daily dosage should be given as two divided doses.

Epilepsy Add-on therapy

Adults

Titration for add-on therapy should begin at 25 to 50 mg as a single (nightly) or divided dose for one week or longer. The dosage should then be increased by 25 to 100 mg/day at weekly or longer intervals to the target dose of 200 to 400 mg/day. The maximum recommended dose should not exceed 1000 mg/day. The daily dosage should be taken as two divided doses.

Children (2 years and over)

Titration for add-on therapy should begin at 1 to 3 mg/kg/day up to 25 mg/day as a single (nightly) dose for the first week. The dosage should then be increased by 1 to 3 mg/kg/day at weekly or

longer intervals to the recommended total daily dose of 5 to 9 mg/kg/day. Daily doses up to 30 mg/kg have been studied and were generally well tolerated. The daily dosage should be given as two divided doses.

		Monotherapy	Add-on therapy
Adults	Starting dose	25 mg as a single (nightly) dose for one week (or longer).	25 to 50 mg as a single (nightly) or divided dose for one week (or longer).
	Escalation dose	Increase by 25 to 50 mg/day at weekly or longer intervals.	Increase by 25 to 100 mg/day at weekly or longer intervals.
	Target dose	100 mg/day	200 to 400 mg/day
	Maximum dose	Up to 500 mg/day ¹	Up to 1000 mg/day
Children 2 years & over	Starting dose	0.5 to 1 mg/kg as a single (nightly) dose for the first week.	1 to 3 mg/kg/day up to 25 mg/day as a single (nightly) dose for the first week.
	Escalation dose	Increase by 0.5 to 1 mg/kg/day at weekly or longer intervals.	Increase by 1 to 3 mg/kg/day at weekly or longer intervals.
	Target dose	3 to 6 mg/kg/day	5 to 9 mg/kg/day
	Maximum dose	Up to 500 mg/day	Up to 30 mg/kg/day
<i>Note:</i> Daily doses greater or equal to 50 mg should be taken as two divided doses.			
¹ Some patients with refractory epilepsy have tolerated doses of 1000 mg/day.			

It is not necessary to monitor topiramate plasma concentrations to optimise TAMATE therapy. For patients receiving concomitant phenytoin and carbamazepine, dosage adjustment for TAMATE may be required (see **SECTION 4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS**).

Migraine

Adults

Titration should begin at 25 mg nightly for 1 week. The dosage should then be increased weekly in increments of 25 mg/day. If the patient is unable to tolerate the titration regimen, longer intervals between dose adjustments can be used.

The recommended total daily dose of TAMATE as treatment for prophylaxis of migraine headache is 100 mg/day administered in two divided doses. Some patients may experience a benefit at a total daily dose of 50 mg/day. Patients have received a total daily dose up to 200 mg/day. Dose and titration should be guided by clinical outcome.

Use in the elderly

Caution is advised during titration in the elderly with renal disease and/or hepatic impairment (see **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE**).

Use in patients with hepatic and/or renal impairment

Caution is advised during titration in patients with renal disease and/or hepatic impairment (see **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE**). Patients with moderate and severe renal impairment may require a dose reduction. Half of the usual starting and maintenance dose is recommended (see **section 5.2 Pharmacokinetic Properties**).

Use in patients undergoing haemodialysis

Topiramate is cleared by haemodialysis. To avoid rapid reduction in topiramate plasma concentration during haemodialysis, a supplemental dose of TAMATE equal to approximately one-half the daily dose should be administered on haemodialysis days. The supplemental dose

should be administered in divided doses at the beginning and completion of the haemodialysis procedure. The supplemental dose may differ based on the characteristics of the dialysis equipment being used (see **section 5.2 Pharmacokinetic Properties**). The actual adjustment should take into account:

- the duration of the dialysis,
- the clearance rate of the dialysis system being used, and
- the effective renal clearance of topiramate in the patient being dialysed.

Drug withdrawal and Dosage reduction

In patients with or without a history of seizures or epilepsy, antiepileptic drugs, including TAMATE, should be gradually withdrawn to minimize the potential for seizures or of increased seizure frequency. In situations where rapid withdrawal of TAMATE is medically required, appropriate monitoring is recommended.

4.3 CONTRAINDICATIONS

Hypersensitivity to any component of this product.

4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Withdrawal of Topiramate

In patients with or without a history of seizures or epilepsy, antiepileptic medicines including topiramate should be gradually withdrawn to minimise the potential for seizures or increased seizure frequency. In clinical trials, daily dosages were decreased in weekly intervals by 50 – 100mg in adults receiving topiramate at doses up to 100mg/day for migraine prophylaxis. In clinical trials of children, topiramate was gradually withdrawn over a 2 – 8 week period. In situations where rapid withdrawal of topiramate is medically required, appropriate monitoring is recommended.

Topiramate has not been studied in patients with a history of psychiatric disorders. Given the reported association of certain antiepileptic agents and psychiatric disturbances, topiramate should be used with caution in patients with a prior psychiatric history.

Identified Precautions

Hydration

Maintaining adequate hydration whilst using topiramate is very important. Hydration can reduce the risk of nephrolithiasis. Proper hydration prior to and during activities such as exercise or exposure to warm temperatures may reduce the risk of heat-related adverse events.

Suicidal Behaviour and Ideation

Antiepileptic drugs (AEDs), including topiramate, increase the risk of suicidal thoughts or behaviour in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behaviour, and/or any unusual changes in mood or behaviour.

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% CI:1.2, 2.7) of suicidal thinking or behaviour compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behaviour or ideation among 27,863 AED-treated patients was 0.43%, compared to 0.24% among 16,029 placebo-treated patients, representing an increase of approximately one case of suicidal thinking or behaviour for every 530 patients treated. There were four suicides in drug-treated patients in

the trials and none in placebo-treated patients, but the number is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behaviour with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behaviour beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behaviour was generally consistent among drugs in the data analysed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5 to 100 years) in the clinical trials analysed.

Table 2 shows absolute and relative risk by indication for all evaluated AEDs.

Indication	Placebo Patients with Events per 1000 Patients	Drug Patients with Events per 1000 Patients	Relative Risk: Incidence of Events in Drug Patients/Incidence in Placebo Patients	Risk Difference: Additional Drug Patients with Events per 1000 Patients
Epilepsy	1.0	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behaviour was higher in clinical trials for epilepsy than in clinical trials for psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.

In double-blind clinical trials with topiramate in approved and investigational indications, suicide related events (suicidal ideation, suicide attempts, and suicide) occurred at a frequency of 0.5% in topiramate treated patients (46 out of 8,652 patients treated) compared to 0.2% treated with placebo (8 out of 4,045 patients treated). One completed suicide was reported in a bipolar disorder double-blind trial in a patient on topiramate.

Anyone considering prescribing topiramate or any other AED must balance the risk of suicidal thoughts or behaviour with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behaviour. Should suicidal thoughts and behaviour emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

Patients, and, when appropriate, their caregivers, and families should be informed that AEDs increase the risk of suicidal thoughts and behaviour and should be advised of the need to be alert for the emergence or worsening of the signs and symptoms of depression, any unusual changes in mood or behaviour or the emergence of suicidal thoughts, behaviour or thoughts about self-harm. Behaviours of concern should be reported immediately to the treating doctor.

Nephrolithiasis

Patients, especially those with a predisposition to nephrolithiasis, may be at increased risk for renal stone formation (none of 216 placebo versus 1.6% of 1446 patients who had received topiramate were reported to have nephrolithiasis) and associated signs and symptoms such as renal colic, renal pain or flank pain.

Risk factors for nephrolithiasis include prior stone formation, a family history of nephrolithiasis and hypercalciuria (see **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE, Metabolic Acidosis**), and gender (male). None of these risk factors can reliably predict stone formation during topiramate treatment. In addition, patients taking other medication associated with nephrolithiasis may be at increased risk.

Oligohydrosis and Hyperthermia

Oligohydrosis (decreased sweating) and anhidrosis, infrequently resulting in hospitalization, has been reported in association with topiramate use. Decreased sweating and an elevation in body temperature above normal characterized these cases. Some of the cases were reported after exposure to elevated environmental temperature.

The majority of the reports have been in children. Patients, especially paediatric patients, treated with topiramate should be monitored closely for evidence of decreased sweating and increased body temperature, especially in hot weather. Caution should be used when topiramate is prescribed with other drugs that predispose patients to heat-related disorders; these drugs include, but are not limited to, other carbonic anhydrase inhibitors and drugs with anticholinergic activity.

Patients, especially paediatric patients, treated with topiramate should be monitored closely for evidence of decreased sweating and increased body temperature, especially in hot weather.

Use in patients with renal impairment

The major route of elimination of unchanged topiramate and its metabolites is via the kidney. Renal elimination is dependent on renal function and is independent of age. Patients with moderate or severe renal impairment may take 10 to 15 days to reach steady-state plasma concentrations as compared to 4 to 8 days in patients with normal renal function.

In all patients the titration schedule should be guided by clinical outcome (i.e. seizure control, avoidance of side effects) and the knowledge that subjects with known renal impairment may require a longer time to reach steady state at each dose.

Use in patients with hepatic impairment

In hepatically-impaired patients, topiramate should be administered with caution as the clearance of topiramate may be decreased.

Acute myopia and secondary angle closure glaucoma

A syndrome consisting of acute myopia associated with secondary angle closure glaucoma has been reported in patients receiving topiramate. Symptoms include acute onset of decreased visual acuity and/or ocular pain. Ophthalmological findings can include myopia, anterior chamber shallowing, ocular hyperaemia (redness) and increased intraocular pressure.

Mydriasis may or may not be present. This syndrome may be associated with supraciliary effusion resulting in anterior displacement of the lens and iris, with secondary angle closure glaucoma. Symptoms typically occur within 1 month of initiating topiramate therapy. In contrast to primary narrow angle glaucoma, which is rare under 40 years of age, secondary angle closure glaucoma associated with topiramate has been reported in paediatric patients as well as adults. Treatment includes discontinuation of topiramate as rapidly as possible in the judgement of the treating physician and appropriate measures to reduce intraocular pressure. These measures generally result in a decrease in intraocular pressure.

Elevated intraocular pressure of any aetiology, if left untreated, can lead to serious sequelae including permanent vision loss.

Visual field defects

Visual field defects have been reported in patients receiving topiramate independent of elevated intraocular pressure. In clinical trials most of these events were reversible after topiramate discontinuation however, some cases were not. In a large proportion of post marketing case reports reversibility was unknown, but

in cases where an outcome was reported, the majority were reversible. If visual problems occur at any time during topiramate treatment, consideration should be given to discontinuing the drug.

Metabolic Acidosis

Hyperchloremic, non-anion gap, metabolic acidosis (i.e. decreased serum bicarbonate below the normal reference range in the absence of respiratory alkalosis) is associated with topiramate treatment. This decrease in serum bicarbonate is due to the inhibitory effect of topiramate on renal carbonic anhydrase. Generally, the decrease in bicarbonate occurs early in treatment although it can occur at any time during treatment. These decreases are usually mild to moderate (average decrease of 4mmol/L as doses of 100mg/day or above in adults and at approximately 6mg/kg/day in paediatric patients).

Rarely, patients have experienced decreases to values below 10mmol/L. Conditions or therapies that predispose to acidosis (such as renal disease, severe respiratory disorders, status epilepticus, diarrhoea, surgery, ketogenic diet, or certain medicines) may be additive to the bicarbonate lowering effects of topiramate.

Chronic, untreated metabolic acidosis may increase the risk of nephrolithiasis or nephrocalcinosis (see **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE, Nephrolithiasis**).

In adults, the incidence of persistent treatment-emergent decreases in serum bicarbonate (levels of <20mmol/L at two consecutive visits or at the final visit) in controlled clinical trials for adjunctive treatment of epilepsy was 32% for 400mg/day, and 1% for placebo. Metabolic acidosis has been observed at doses as low as 50mg/day. The incidence of a markedly abnormally low serum bicarbonate (i.e. absolute value <17mmol/L and >5mmol/L decrease from pre-treatment) in these trials was 3% for 400mg/day, and 0% for placebo. Serum bicarbonate levels have not been systematically evaluated at daily doses greater than 400mg/day.

The incidence of persistent treatment-emergent decreases in serum bicarbonate in placebo-controlled trials for adults for prophylaxis of migraine was 44% for 200mg/day, 39% for 100mg/day, 23% for 50mg/day, and 7% for placebo. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17mmol/L and >5mmol/L decrease from pre-treatment) in these trials was 11% for 200mg/day, 9% for 100mg/day, 2% for 50mg/day, and <1% for placebo.

In paediatric patients (<16 years of age), the incidence of persistent treatment-emergent decreases in serum bicarbonate in placebo-controlled trials for adjunctive treatment of Lennox-Gastaut syndrome or refractory partial onset seizures was 67% for topiramate (at approximately 6mg/kg/day), and 10% for placebo. The incidence of a markedly abnormally low serum bicarbonate (i.e., absolute value <17mmol/L and >5mmol/L decrease from pre-treatment) in these trials was 11% for topiramate and 0% for placebo. Cases of moderately severe metabolic acidosis have been reported in patients as young as 5 months old, especially at daily doses above 5mg/kg/day.

Some manifestations of acute or chronic metabolic acidosis may include hyperventilation, nonspecific symptoms such as fatigue and anorexia, or more severe sequelae including cardiac arrhythmias or stupor. Chronic, untreated metabolic acidosis may increase the risk for nephrolithiasis or nephrocalcinosis, and may also result in osteomalacia (referred to as rickets in pediatric patients) and/or osteoporosis with an increased risk for fractures. Chronic metabolic acidosis in pediatric patients can reduce growth rates. A reduction in growth rate may eventually decrease the maximal height achieved. The effect of topiramate on growth and bone-related sequelae has not been systematically investigated in pediatric or adult populations.

Depending on underlying conditions, appropriate evaluation including serum bicarbonate levels is recommended with topiramate therapy. If metabolic acidosis develops and persists, consideration should be given to reducing the dose or discontinuing topiramate (by using dose tapering).

Hyperammonaemia and encephalopathy

Hyperammonemia with or without encephalopathy has been reported with topiramate treatment (see **section 4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)**). The risk for hyperammonemia with

topiramate appears dose-related. Hyperammonemia has been reported more frequently when topiramate is used concomitantly with valproic acid (see **section 4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS**).

Clinical symptoms of hyperammonemic encephalopathy often include acute alterations in level of consciousness and/or cognitive function with lethargy. In most cases, hyperammonemic encephalopathy abated with discontinuation of treatment. In patients who develop unexplained lethargy, or changes in mental status associated with topiramate monotherapy or adjunctive therapy, it is recommended to consider hyperammonemic encephalopathy and measuring ammonia levels.

Mood Disturbances/Depression

An increased incidence of mood disturbances and depression has been observed during topiramate treatment. Psychiatric/behavioural disturbances (depression or mood problems) in the majority of affected patients were dose related for both the add-on epilepsy and migraine populations.

Women of childbearing potential

Topiramate may cause foetal harm when administered to a pregnant woman. There is an increased risk of pre-term labour, premature delivery and congenital malformations associated with the use of AEDs, including topiramate.

Topiramate should be used during pregnancy only if the potential benefit justifies the potential risk to the foetus (see **section 4.6 FERTILITY, PREGNANCY and LACTATION**).

Use in the Elderly

See **section 4.2 DOSE AND METHOD OF ADMINISTRATION - Use in the elderly**.

Paediatric Use

See **section 4.2 DOSE AND METHOD OF ADMINISTRATION – Children (2 years and over)**.

Effects on Laboratory Tests

No data available.

4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS

Studies in mice receiving concomitant administration of topiramate and carbamazepine or phenobarbitone showed synergistic anticonvulsant activity, while combination with phenytoin showed additive anticonvulsant activity.

Effects of topiramate on other antiepileptic drugs

The addition of topiramate to other antiepileptic drugs (phenytoin, carbamazepine, valproic acid, phenobarbitone, primidone) has no effect on their steady-state plasma concentrations, except in the occasional patient, where the addition of topiramate to phenytoin may result in an increase of plasma concentrations of phenytoin. This is possibly due to inhibition of a specific enzyme polymorphic isoform (CYP2C19). Consequently, any patient on phenytoin showing clinical signs or symptoms of toxicity should have phenytoin levels monitored.

Effects of other antiepileptic drugs on topiramate

The metabolic breakdown of topiramate is increased in patients receiving concomitant antiepileptic therapy with agents that are inducers of drug metabolising enzymes. The increased metabolic breakdown results in up to 1.5 times higher clearance of topiramate.

Phenytoin and carbamazepine decrease the plasma concentration of topiramate. The addition or withdrawal of phenytoin or carbamazepine to topiramate therapy may require an adjustment in dosage of the latter. This should be done by titrating to clinical effect.

The addition or withdrawal of valproic acid does not produce clinically significant changes in plasma concentrations of topiramate and, therefore, does not warrant dosage adjustment of topiramate.

The results of these interactions are summarised in **Table 3**.

Table 3: Summary of AED interactions with topiramate		
AED Co-administered	AED Concentration	Topiramate Concentration
Phenytoin	↔ *	↓** (48%)
Carbamazepine (CBZ)	↔	↓ (40%)
Valproic Acid	↔	↔
Phenobarbitone	↔	N
Primidone	↔	N
Lamotrigine	↔	↔
↔ = No effect on plasma concentration * = Plasma concentrations increase in occasional patients *↓ = Plasma concentrations decrease ** = Approximately 35% decrease in plasma C _{maxss} and 57% decrease in plasma C _{minss} concentrations N = Not studied AED = antiepileptic drug NOTE: No data are available on the use of topiramate with lamotrigine, gabapentin or vigabatrin.		

No data are available on the use of topiramate with vigabatrin.

Other drug interactions

Digoxin

In a single-dose study, serum digoxin area under plasma concentration curve (AUC) decreased 12% due to concomitant administration of topiramate. The clinical relevance of this observation has not been established. When topiramate is added or withdrawn in patients on digoxin therapy, careful attention should be given to the routine monitoring of serum digoxin.

CNS Depressants

Concomitant administration of topiramate and alcohol or other CNS depressant drugs has not been evaluated in clinical studies. It is recommended that topiramate not be used concomitantly with alcohol or other CNS depressant drugs.

Oral Contraceptives

In a pharmacokinetic interaction study in healthy volunteers with a concomitantly administered combination oral contraceptive product containing 1 mg norethisterone (NET) plus 35 mcg ethinyl oestradiol (EO), topiramate given in the absence of other medications at doses of 50 to 200mg/day was not associated with statistically significant changes in mean exposure (AUC) to either component of the oral contraceptive. In another study, exposure to EO was statistically significantly decreased at doses of 200, 400, and 800mg/day (18%, 21%, and 30%, respectively) when given as adjunctive therapy in patients taking valproic acid. In both studies, topiramate (50mg/day to 800mg/day) did not significantly effect exposure to NET. Although there was a dose dependent decrease in EO exposure for doses between 200-800mg/day, there was no significant dose dependent change in EO exposure for doses of 50-200mg/day.

The clinical significance of the changes observed is not known. The possibility of decreased contraceptive efficacy and increased breakthrough bleeding should be considered in patients taking combination oral contraceptive products with topiramate. Patients taking oestrogen containing contraceptives should be

asked to report any change in their bleeding patterns. Contraceptive efficacy can be decreased even in the absence of breakthrough bleeding.

Lithium

In healthy volunteers, there was an observed reduction (18% for AUC) in systemic exposure for lithium during concomitant administration with topiramate 200mg/day. In patients with bipolar disorder, the pharmacokinetics of lithium were unaffected during treatment with topiramate at doses of 200mg/day; however, there was an observed increase in systemic exposure (26% for AUC) following topiramate doses of up to 600mg/day. Lithium levels should be monitored when co-administered with topiramate.

Risperidone

Drug-drug interaction studies conducted under single and multiple dose conditions in healthy volunteers and patients with bipolar disorder yielded similar results. When administered concomitantly with topiramate at escalating doses of 100, 250 and 400mg/day there was a reduction in risperidone (administered at doses ranging from 1 to 6mg/day) systemic exposure (16% and 33% for steady-state AUC at the 250 and 400mg/day doses, respectively). Minimal alterations in the pharmacokinetics of the total active moiety (risperidone plus 9-hydroxyrisperidone) and no alterations for 9-hydroxyrisperidone were observed. The clinical relevance of the observed, apparently not statistically significant changes in the systemic exposure of the total active moiety (risperidone plus 9-hydroxyrisperidone) or of topiramate is not known.

Hydrochlorothiazide (HCTZ)

A drug-drug interaction study conducted in healthy volunteers evaluated the steady-state pharmacokinetics of HCTZ (25 mg q24h) and topiramate (96 mg q12h) when administered alone and concomitantly. The results of this study indicate that topiramate C_{max} increased by 27% and AUC increased by 29% when HCTZ was added to topiramate. The clinical significance of this change is unknown. The addition of HCTZ to topiramate therapy may require an adjustment of the topiramate dose. Clinical laboratory results indicated decreases in serum potassium after topiramate or HCTZ administration, which were greater when HCTZ and topiramate were administered in combination.

Metformin

A drug-drug interaction study conducted in healthy volunteers evaluated the steady-state pharmacokinetics of metformin and topiramate in plasma when metformin was given alone and when metformin and topiramate were given simultaneously. The results of this study indicated that metformin mean C_{max} and mean AUC_{0-12h} increased by 18% and 25%, respectively, while mean CL/F decreased 20% when metformin was co-administered with topiramate. Topiramate did not affect metformin t_{max} . The clinical significance of the effect of topiramate on metformin pharmacokinetics is unclear. Oral plasma clearance of topiramate appears to be reduced when administered with metformin. The extent of change in the clearance is unknown. The clinical significance of the effect of metformin on topiramate pharmacokinetics is unclear. When topiramate is added or withdrawn in patients on metformin therapy, careful attention should be given to the routine monitoring for adequate control of their diabetic disease state.

Pioglitazone

A drug-drug interaction study conducted in healthy volunteers evaluated the steady-state pharmacokinetics of topiramate and pioglitazone when administered alone and concomitantly. A 15% decrease in the $AUC_{t,ss}$ of pioglitazone with no alteration in $C_{max,ss}$ was observed. This finding was not statistically significant. In addition, a 13% and 16% decrease in $C_{max,ss}$ and $AUC_{t,ss}$ respectively, of the active hydroxy-metabolite was noted as well as a 60% decrease in $C_{max,ss}$ and $AUC_{t,ss}$ of the active keto-metabolite. The clinical significance of these findings is not known. When topiramate is added to pioglitazone therapy or pioglitazone is added to topiramate therapy, careful attention should be given to the routine monitoring of patients for adequate control of their diabetic disease state.

Glibenclamide

A drug-drug interaction study conducted in patients with type 2 diabetes evaluated the steady-state pharmacokinetics of glibenclamide (5mg/day) alone and concomitantly with topiramate (150mg/day). There was a 25% reduction in glibenclamide AUC_{24} during topiramate administration. Systemic exposure

of the active metabolites, 4-trans-hydroxy-glibenclamide (M1) and 3-cis-hydroxyglibenclamide (M2), were also reduced by 13% and 15%, respectively. The steady-state pharmacokinetics of topiramate were unaffected by concomitant administration of glibenclamide. When topiramate is added to glibenclamide therapy or glibenclamide is added to topiramate therapy, careful attention should be given to the routine monitoring of patients for adequate control of their diabetic disease state.

Other forms of interactions

Agents predisposing to nephrolithiasis

Topiramate, when used concomitantly with other agents predisposing to nephrolithiasis, may increase the risk of nephrolithiasis. While using topiramate, agents like these should be avoided since they may create a physiological environment that increases the risk of renal stone formation.

Valproic Acid

Concomitant administration of topiramate and valproic acid has been associated with hyperammonemia with or without encephalopathy in patients who have tolerated either drug alone. In most cases, symptoms and signs abated with discontinuation of either drug (see **section 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE and section 4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)**). This adverse effect is not due to a pharmacokinetic interaction.

Hypothermia, defined as an unintentional drop in body core temperature to $<35^{\circ}\text{C}$, has been reported in association with concomitant use of topiramate and valproic acid (VPA) both in conjunction with hyperammonemia and in the absence of hyperammonemia. This adverse event in patients using concomitant topiramate and valproate can occur after starting topiramate treatment or after increasing the daily dose of topiramate.

Vitamin K-antagonist anticoagulant medications

Decreased Prothrombin Time/International Normalized Ratio (PT/INR) responses have been reported following concomitant administration of topiramate with vitamin K-antagonist anticoagulant medications. Closely monitor INR during concomitant administration of topiramate therapy with vitamin K-antagonist anticoagulant medications.

Additional Pharmacokinetic Drug Interaction Studies.

Clinical studies have been conducted to assess the potential pharmacokinetic drug interaction between topiramate and other agents. The changes in C_{max} or AUC as a result of the interactions are summarized below in **Table 4**. The second column (concomitant drug concentration) describes what happens to the concentration of the concomitant drug listed in the first column when topiramate is added. The third column (topiramate concentration) describes how the coadministration of a drug listed in the first column modifies the concentration of topiramate.

Concomitant Drug	Concomitant Drug Concentration	Topiramate Concentration
Amitriptyline	↔ 20% increase in C_{max} and AUC of nortriptyline metabolite	NS
Dihydroergotamine (Oral and Subcutaneous)	↔	↔
Haloperidol	↔ 31% increase in AUC of the reduced metabolite	NS
Propranolol	↔ 17% Increase in C_{max} for 4-OH propranolol (TPM 50mg q12h)	9% and 16% Increase in C_{max} , 9% and 17% increase in AUC (40 mg and 80mg propranolol q12h respectively)
Sumatriptan (Oral and Subcutaneous)	↔	NS

Concomitant Drug	Concomitant Drug Concentration	Topiramate Concentration
Pizotifen	↔	↔
Diltiazem	25% decrease in AUC of diltiazem and 18% decrease in DEA, and ↔ for DEM	20% increase in AUC
*Venlafaxine	↔	↔
*Flunarizine	16% increase in AUC (TPM 50mg q12h) ^b	↔

↔ = No effect on C_{max} and AUC (≤ 15% change) of the parent compound
^b = Flunarizine AUC increased 14% in subjects taking flunarizine alone. Increase in exposure may be attributed to accumulation during achievement of steady state.
 NS = Not studied
 *DEA = des acetyl diltiazem, DEM = N-demethyl diltiazem

4.6 FERTILITY, PREGNANCY AND LACTATION

Effects on Fertility

There were no effects on fertility or reproductive parameters in rats following oral administration of topiramate at doses up to 100mg/kg/day, with estimated exposures (plasma AUC) less than human exposure at the maximal recommended clinical dose. Oral administration of topiramate to juvenile rats did not affect subsequent reproductive development, mating or fertility (see **4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE - Growth and Development**).

Use in Pregnancy

Pregnancy Category: D

When administered orally during organogenesis, topiramate was teratogenic in mice, rats and rabbits at maternal exposures (plasma AUC) less than clinical exposure at the maximal recommended dose. In mice, the numbers of foetal malformations (primarily craniofacial abnormalities) were increased at all dose levels tested. The malformations in rats (limb reduction defects) and rabbits (axial and costal skeletal defects) were similar to those seen with carbonic anhydrase inhibitors in these species. Carbonic anhydrase inhibitors have not been associated with malformations in human beings.

There are no studies using topiramate in pregnant women. In post-marketing experience, cases of hypospadias have been reported in male infants exposed in-utero to topiramate, with or without other anticonvulsants. A causal relationship with topiramate has not been established.

There are no adequate and well-controlled studies using topiramate in pregnant women.

Topiramate can cause foetal harm when administered to a pregnant woman. Data from pregnancy registries indicate that infants exposed to topiramate in utero have an increased risk of congenital malformations (e.g., craniofacial defects, such as cleft lip/palate, hypospadias, and anomalies involving various body systems). This has been reported with topiramate monotherapy and topiramate as part of a polytherapy regimen.

Data from the North American AED (NAAED) Pregnancy Registry indicate an increased risk of oral clefts in infants exposed to topiramate monotherapy during the first trimester of pregnancy. The prevalence of oral clefts was 1.4% compared to a prevalence of 0.38% - 0.55% in infants exposed to other AEDs, and a prevalence of 0.07 % in infants of mothers without epilepsy or treatment with other AEDs. The relative risk of oral clefts in topiramate-exposed pregnancies in the NAAED Pregnancy Registry was 21.3 (95% Confidence Interval 7.9 – 57.1) as compared to the risk in a background population of untreated women. The UK Epilepsy and Pregnancy Register reported a similarly increased prevalence of oral clefts of 3.2% among infants exposed to topiramate monotherapy. The observed rate of oral clefts was 16 times higher than the background rate in the UK, which is approximately 0.2%.

In addition, data from these registries and other studies suggest that, compared with monotherapy, there may be an increased risk of teratogenic effects associated with the use of anti-epileptic drugs in combination therapy. The risk has been observed in all doses and effects were reported to be dose-dependent. In women treated with topiramate who have had a child with a congenital malformation, there appears to be an increased risk of malformations in subsequent pregnancies when exposed to topiramate. There is an increased risk of pre-term labour and premature delivery associated with the use of AEDs, including topiramate.

Compared with a reference group not taking antiepileptic drugs, registry data for topiramate monotherapy showed a higher prevalence of low birth weight (<2500 grams). One pregnancy registry reported an increased frequency of infants who were small for gestational age (SGA; defined as birth weight below the 10th percentile corrected for their gestational age, stratified by sex) among those exposed to topiramate monotherapy *in utero*. SGA has been observed in all doses and is dose-dependent. The prevalence of SGA is greater in women who received higher doses of topiramate during pregnancy. In addition, the prevalence of SGA for women who continued topiramate use later in pregnancy is higher compared to women who stopped its use before the third trimester. The long-term consequences of the SGA findings could not be determined. A causal relationship for low birth weight and SGA has not been established. Topiramate should be used during pregnancy only if potential benefit justifies the potential risk to the foetus. In treating and counselling women of childbearing potential, the prescribing physician should weigh the benefits of therapy against the risks. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the foetus.

The risk of having an abnormal child as a result of antiepileptic medication is far outweighed by the danger to the mother and foetus of uncontrolled epilepsy.

It is recommended that:

- Women on antiepileptic drugs (AEDs) receive pregnancy counselling with regard to the risk of foetal abnormalities;
- AEDs should be continued during pregnancy and monotherapy should be used if possible at the lowest effective dose as risk of abnormality is greater in women taking combined medication;
- Folic acid supplementation (5mg) should be commenced four weeks prior to and continue for twelve weeks after conception;
- Specialist prenatal diagnosis including detailed mid-trimester ultrasound should be offered.

Use in Lactation

Radioactivity was detected in milk following oral administration of radiolabelled topiramate to lactating rats. About 1.5% of the dose was recovered in milk in 24 hours, and milk and maternal plasma radioactivity concentrations were similar. The excretion of topiramate has not been evaluated in controlled studies. Limited observation in patients suggests an extensive excretion of topiramate in breast milk. Lactating women should be advised not to breastfeed during treatment with topiramate.

Growth and Development

In juvenile rats, oral administration of topiramate at doses up to 300 mg/kg/day during the period of development corresponding to infancy, childhood, and adolescence resulted in toxicities similar to those in adult animals (decreased food consumption with decreased body weight gain, centrolobullar hepatocellular hypertrophy and slight urothelial hyperplasia in the urinary bladder). There were no relevant effects on long bone (tibia) growth or bone (femur) mineral density, preweaning and reproductive development, neurological development (including assessments on memory and learning), mating and fertility or hysterotomy parameters. Exposure (plasma AUC) was up to 2-fold human exposure at the maximal recommended clinical dose.

4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

Topiramate acts on the central nervous system and may produce drowsiness, dizziness or other related symptoms. It may also cause visual disturbances and/ or blurred vision. These adverse events are potentially dangerous in patients driving a vehicle or operating machinery, particularly until the individual patient's experience with the drug is established.

4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)

Clinical Trial Data

The safety of topiramate was evaluated from a clinical trial database consisting of 4111 patients (3182 on topiramate and 929 on placebo) who participated in 20 double-blind trials and 2847 patients who participated in 34 open-label trials, respectively, for the treatment of primary generalized tonic-clonic seizures, partial onset seizures, seizures associated with Lennox-Gastaut syndrome, newly or recently diagnosed epilepsy or migraine. The information presented in this section was derived from pooled data.

The majority of all adverse reactions were mild to moderate in severity.

Increased risk for Bleeding

Topiramate treatment is associated with an increased risk for bleeding. In a pooled analysis of placebo-controlled studies of approved and unapproved indications, bleeding was more frequently reported as an adverse event for topiramate than for placebo (4.5% versus 3.0% in adult patients, and 4.4% versus 2.3% in paediatric patients). In this analysis, the incidence of serious bleeding events for topiramate and placebo was 0.3% versus 0.2% for adult patients, and 0.4% versus 0% for paediatric patients.

Adverse bleeding reactions reported with topiramate ranged from mild epistaxis, ecchymosis, and increased menstrual bleeding to life-threatening haemorrhages. In patients with serious bleeding events, conditions that increased the risk for bleeding were often present, or patients were often taking drugs that cause thrombocytopenia (other antiepileptic drugs) or affect platelet function or coagulation (e.g., aspirin, non-steroidal anti-inflammatory drugs, selective serotonin reuptake inhibitors, or warfarin or other anticoagulants).

Double-Blind, Placebo-Controlled Data, Adjunctive Epilepsy Trials – Adult Patients

Adverse Drug Reactions (ADRs) reported in $\geq 1\%$ of topiramate-treated adult patients in double-blind, placebo-controlled adjunctive epilepsy trials are shown in **Table 5**. ADRs that had an incidence $>5\%$ in the recommended dose range (200 to 400 mg/day) in adults in double-blind, placebo-controlled adjunctive epilepsy studies in descending order of frequency included somnolence, dizziness, fatigue, irritability, weight decreased, bradyphrenia, paresthesias, diplopia, coordination abnormal, nausea, nystagmus, lethargy, anorexia, dysarthria, vision blurred, decreased appetite, memory impairment and diarrhoea.

Table 5: Adverse Drug Reactions Reported by $\geq 1\%$ of Topiramate-Treated Adult Patients in Double-Blind, Placebo-Controlled, Adjunctive Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate 200 – 400 mg/day (N=354) %	Topiramate 600 – 1000 mg/day (N=437) %	Placebo (N=382) %
Metabolism and Nutrition Disorders			
Anorexia	5.4	6.2	1.8
Decreased appetite	5.1	8.7	3.7
Psychiatric Disorders			
Bradyphrenia	8.2	19.5	3.1
Expressive language disorder	4.5	9.4	1.6
Confusional state	3.1	5.0	0.8
Depression	3.1	11.7	3.4

Table 5: Adverse Drug Reactions Reported by $\geq 1\%$ of Topiramate-Treated Adult Patients in Double-Blind, Placebo-Controlled, Adjunctive Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate 200 – 400 mg/day (N=354) %	Topiramate 600 – 1000 mg/day (N=437) %	Placebo (N=382) %
Insomnia	3.1	6.4	4.5
Aggression	2.8	3.2	1.8
Agitation	1.7	2.3	1.3
Anger	1.7	2.1	0.5
Anxiety	1.7	6.6	2.9
Disorientation	1.7	3.2	1.0
Mood altered	1.7	4.6	1.0
Nervous System Disorders			
Somnolence	17.8	17.4	8.4
Dizziness	16.4	34.1	13.6
Paraesthesia	8.2	17.2	3.7
Coordination abnormal	7.1	11.4	4.2
Nystagmus	6.2	11.7	6.8
Lethargy	5.6	8.0	2.1
Dysarthria	5.4	6.2	1.0
Memory impairment	5.1	10.8	1.8
Disturbance in attention	4.5	11.9	1.8
Tremor	4.0	9.4	5.0
Amnesia	3.4	5.3	1.0
Balance disorder	3.4	3.9	2.4
Hypoaesthesia	3.1	5.9	1.0
Intention tremor	3.1	4.8	2.9
Dysgeusia	1.4	4.3	0.8
Mental impairment	1.4	5.0	1.3
Speech disorder	1.1	2.7	0.5
Eye Disorders			
Diplopia	7.3	12.1	5.0
Vision blurred	5.4	8.9	2.4
Visual disturbance	2.0	1.4	0.3
Gastrointestinal Disorders			
Nausea	6.8	15.1	8.4
Diarrhoea	5.1	14.0	5.2
Abdominal pain upper	3.7	3.9	2.1
Constipation	3.7	3.2	1.8
Stomach discomfort	3.1	3.2	1.3
Dyspepsia	2.3	3.0	2.1
Dry mouth	1.7	3.7	0.3
Abdominal pain	1.1	2.7	0.8
Musculoskeletal and Connective Tissue Disorders			
Myalgia	2.0	2.5	1.3
Muscle spasms	1.7	2.1	0.8
Musculoskeletal chest pain	1.1	1.8	0.3

Table 5: Adverse Drug Reactions Reported by $\geq 1\%$ of Topiramate-Treated Adult Patients in Double-Blind, Placebo-Controlled, Adjunctive Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate 200 – 400 mg/day (N=354) %	Topiramate 600 – 1000 mg/day (N=437) %	Placebo (N=382) %
General Disorders and Administration Site Conditions			
Fatigue	13.0	30.7	11.8
Irritability	9.3	14.6	3.7
Asthenia	3.4	3.0	1.8
Gait disturbance	1.4	2.5	1.3
Investigations			
Weight decreased	9.0	11.9	4.2

Adverse drug reactions reported by $<1\%$ of topiramate-treated adult patients in double-blind, placebo-controlled, adjunctive epilepsy trials included increased appetite, abnormal behaviour, apathy, depressed mood, distractibility, disturbance in sexual arousal, dysphemia, euphoric mood, flat affect, lack of spontaneous speech, mood swings, panic disorder, paranoia, reading disorder, sleep disorder, suicidal ideation, thinking abnormal, aphasia, cerebellar syndrome, cognitive disorder, dysaesthesia, dysgraphia, dyskinesia, formication, parosmia, psychomotor skills impaired, repetitive speech, sensory disturbance, sensory loss, stupor, unresponsive to stimuli, lacrimation increased, presbyopia, visual acuity reduced, deafness, deafness unilateral, hearing impaired, tinnitus, sinus bradycardia, dyspnoea, dyspnoea exertional, paranasal sinus hypersecretion, abdominal discomfort, abdominal tenderness, breath odour, flatulence, hypoaesthesia oral, paraesthesia oral, erythema, hypoaesthesia facial, skin odour abnormal, swelling face, calculus urinary, dysuria, haematuria, incontinence, pollakiuria, renal colic, renal pain, urinary, incontinence, erectile dysfunction, sexual dysfunction, feeling abnormal, feeling drunk, tandem gait test abnormal, and white blood cell count decreased. The recommended dose for adjunctive epilepsy therapy in adults is 200-400 mg/day.

Double-Blind, Placebo-Controlled Data, Adjunctive Epilepsy Trials – Pediatric Patients

ADRs reported in $>2\%$ of topiramate-treated pediatric patients (2 to 16 years of age) in double-blind, placebo-controlled adjunctive epilepsy trials are shown in **Table 6**. ADRs that had an incidence $>5\%$ in the recommended dose range (5 to 9 mg/kg/day) in descending order of frequency included decreased appetite, fatigue, somnolence, lethargy irritability, disturbance in attention, weight decreased, aggression, rash, abnormal behaviour, anorexia, balance disorder, and constipation.

Table 6: Adverse Drug Reactions Reported by $\geq 2\%$ of Topiramate-Treated Paediatric Patients in Double-Blind, Placebo-Controlled, Adjunctive Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate (N=104) %	Placebo (N=102) %
Metabolism and Nutrition Disorders		
Decreased appetite	19.2	12.7
Anorexia	5.8	1.0
Psychiatric Disorders		
Aggression	8.7	6.9
Abnormal behaviour	5.8	3.9
Confusional state	2.9	2.0
Mood altered	2.9	2.0
Nervous System Disorders		
Somnolence	15.4	6.9
Lethargy	13.5	8.8
Disturbance in attention	10.6	2.0

Table 6: Adverse Drug Reactions Reported by $\geq 2\%$ of Topiramate-Treated Paediatric Patients in Double-Blind, Placebo-Controlled, Adjunctive Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate (N=104) %	Placebo (N=102) %
Balance disorder	5.8	2.0
Dizziness	4.8	2.9
Memory impairment	3.8	1.0
Respiratory, Thoracic and Mediastinal Disorders		
Epistaxis	4.8	1.0
Gastrointestinal Disorders		
Constipation	5.8	4.9
Skin and Subcutaneous Tissue Disorders		
Rash	6.7	5.9
General Disorders and Administration Site Conditions		
Fatigue	16.3	4.9
Irritability	11.5	8.8
Gait disturbance	4.8	2.0
Investigations		
Weight decreased	9.6	1.0
<p>Adverse drug reactions reported by $<2\%$ of topiramate-treated pediatric patients in double-blind, placebo-controlled, adjunctive epilepsy trials included leukopenia, lymphadenopathy, thrombocytopenia, increased appetite, anger, middle insomnia, perseveration, dysarthria, dysgeusia, paraesthesia, poor quality sleep, syncope, tremor, diplopia, lacrimation increased, vision blurred, sinus bradycardia, paranasal sinus hypersecretion, abdominal pain, flatulence, gastroesophageal reflux disease, glossodynia, alopecia, skin discolouration, musculoskeletal stiffness, incontinence, pollakiuria, asthenia, feeling abnormal, malaise, and thirst.</p> <p>Nausea and headache were not considered ADRs based on case review which indicated that these events could be attributed to other causes, including concomitant use of other medications or an intervening illness</p> <p>The recommended dose for adjunctive epilepsy therapy in children (2-16 years of age) is 5 to 9 mg/kg/day.</p>		

Double-Blind, Controlled Data, Monotherapy Epilepsy Trials – Adult Patients

ADRs reported in $\geq 1\%$ of topiramate-treated adult patients in double-blind, controlled monotherapy epilepsy trials are shown in **Table 7**. ADRs that had an incidence $>5\%$ at the recommended dose (400 mg/day) in descending order of frequency included paraesthesia, weight decreased, fatigue, anorexia, depression, memory impairment, anxiety, diarrhoea, asthenia, dysgeusia, and hypoesthesia.

Table 7: Adverse Drug Reactions Reported by $\geq 1\%$ of Topiramate-Treated Adult Patients in Double-Blind, Controlled Monotherapy Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate 50 mg/day (N=257) %	Topiramate 400 mg/day (N=153) %
Blood and Lymphatic System Disorders		
Anaemia	0.8	2.0
Metabolism and Nutrition Disorders		
Anorexia	3.5	12.4
Decreased appetite	2.3	2.6

Table 7: Adverse Drug Reactions Reported by $\geq 1\%$ of Topiramate-Treated Adult Patients in Double-Blind, Controlled Monotherapy Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate 50 mg/day (N=257) %	Topiramate 400 mg/day (N=153) %
Psychiatric Disorders		
Depression	4.3	8.5
Anxiety	3.9	6.5
Bradyphrenia	2.3	4.6
Expressive language disorder	3.5	4.6
Depressed mood	0.8	2.6
Mood altered	0.4	2.0
Mood swings	1.6	2.0
Nervous System Disorders		
Paraesthesia	18.7	40.5
Memory impairment	1.2	7.2
Dysgeusia	2.3	5.9
Hypoaesthesia	4.3	5.2
Balance disorder	1.6	3.3
Dysarthria	1.6	2.6
Cognitive disorder	0.4	2.0
Lethargy	1.2	2.0
Mental impairment	0.8	2.0
Psychomotor skills impaired	0	2.0
Sedation	0	1.3
Visual field defect	0.4	1.3
Eye Disorders		
Dry eye	0	1.3
Ear and Labyrinth Disorders		
Ear pain	0	1.3
Tinnitus	1.6	1.3
Respiratory, Thoracic and Mediastinal Disorders		
Dyspnoea	1.2	2.0
Rhiorrhoea	0	1.3
Gastrointestinal Disorders		
Diarrhoea	5.4	6.5
Paraesthesia oral	1.2	3.3
Dry mouth	0.4	2.6
Gastritis	0.8	2.6
Abdominal pain	1.2	2.0
Gastroesophageal reflux disease	0.4	2.0
Gingival bleeding	0	1.3
Skin and Subcutaneous Tissue Disorders		
Rash	0.4	3.9
Alopecia	1.6	3.3
Pruritis	0.4	3.3
Hypoaesthesia facial	0.4	2.0

Table 7: Adverse Drug Reactions Reported by $\geq 1\%$ of Topiramate-Treated Adult Patients in Double-Blind, Controlled Monotherapy Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate 50 mg/day (N=257) %	Topiramate 400 mg/day (N=153) %
Pruritis generalised	0	1.3
Musculoskeletal and Connective Tissue Disorders		
Muscle spasms	2.7	3.3
Arthralgia	1.9	2.0
Muscle twitching	0.4	1.3
Renal and Urinary Disorders		
Nephrolithiasis	0	2.6
Dysuria	0.8	2.0
Pollakiuria	0.8	2.0
Reproductive System and Breast Disorders		
Erectile dysfunction	0.8	1.3
General Disorders and Administration Site Conditions		
Fatigue	15.2	14.4
Asthenia	3.5	5.9
Irritability	3.1	3.3
Investigations		
Weight decreased	7.0	17.0
Adverse drug reactions reported by $<1\%$ of topiramate-treated adult patients in double-blind, controlled monotherapy epilepsy trials included lymphadenopathy, increased appetite, polydipsia, agitation, anger, dysphemia, euphoric mood, initial insomnia, suicidal ideation, drooling, hypogeusia, poor quality sleep, sensory disturbance, accommodation disorder, amblyopia, diplopia, palpitations, abdominal discomfort, breath odour, glossodynia, stomach discomfort, anhidrosis, urticaria localised, muscular weakness, and thirst. The recommended dose for monotherapy therapy in adults is 400 mg/day.		

Double-Blind, Controlled Data, Monotherapy Epilepsy Trials – Paediatric Patients

ADRs reported in $\geq 2\%$ of topiramate-treated pediatric patients (10 to 16 years of age) in double-blind, controlled monotherapy epilepsy trials are shown in **Table 8**. ADRs that had an incidence $>5\%$ at the recommended dose (400 mg/day) in descending order of frequency included weight decreased, paraesthesia, diarrhoea, disturbance in attention, pyrexia, and alopecia.

Table 8: Adverse Drug Reactions Reported by $\geq 2\%$ of Topiramate-Treated Paediatric Patients in Double-Blind, Controlled Monotherapy Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate 50 mg/day (N=77) %	Topiramate 400 mg/day (N=63) %
Metabolism and Nutrition Disorders		
Decreased appetite	1.3	4.8
Psychiatric Disorders		
Bradyphrenia	0	4.8
Mood altered	1.3	4.8
Depression	0	3.2

Table 8: Adverse Drug Reactions Reported by $\geq 2\%$ of Topiramate-Treated Paediatric Patients in Double-Blind, Controlled Monotherapy Epilepsy Trials

System/Organ Class Adverse Reaction	Topiramate 50 mg/day (N=77) %	Topiramate 400 mg/day (N=63) %
Nervous System Disorders		
Paraesthesia	3.9	15.9
Disturbance in attention	3.9	7.9
Ear and Labyrinth Disorders		
Vertigo	0	3.2
Respiratory, Thoracic and Mediastinal Disorders		
Epitaxis	0	3.2
Gastrointestinal Disorders		
Diarrhoea	3.9	9.5
Vomiting	3.9	4.8
Skin and Subcutaneous Tissue Disorders		
Alopecia	0	6.3
General Disorders and Administration Site Conditions		
Pyrexia	0	6.3
Asthenia	0	4.8
Investigations		
Weight decreased	7.8	20.6
Social Circumstances		
Learning disability	0	3.2
Adverse drug reactions reported by $<2\%$ of topiramate-treated pediatric patients in double-blind, controlled monotherapy epilepsy trials included eosinophilia, hypersensitivity, increased appetite, abnormal behaviour, apathy, confusional state, crying, distractibility, expressive language disorder, insomnia, mood swings, lethargy, mental impairment, poor quality sleep, diplopia, nasal congestion, rhinorrhoea, abdominal discomfort, gastritis, pruritus, rash, skin discolouration, urticaria, arthralgia, micturition urgency, pollakiuria, feeling abnormal, and hyperthermia.		
The recommended dose for monotherapy therapy in children 10 years and older is 400 mg/day.		

Double-Blind, Placebo-Controlled Data, Migraine Prophylaxis Trials – Adult Patients

ADRs reported in $\geq 1\%$ of topiramate-treated adult patients in double-blind, placebo-controlled migraine prophylaxis trials are shown in **Table 9**. ADRs that had an incidence $>5\%$ at the recommended dose (100 mg/day) in descending order of frequency included paraesthesia, fatigue, nausea, diarrhoea, weight decreased, dysgeusia, anorexia, decreased appetite, insomnia, hypoesthesia, disturbance in attention, anxiety, somnolence, and expressive language disorder.

Table 9: Adverse Drug Reactions Reported by $\geq 1\%$ of Topiramate-Treated Adult Patients in Double-Blind, Placebo-Controlled Migraine Prophylaxis Trials

System/Organ Class Adverse Reaction	Topiramate 50 mg/day (N=227) %	Topiramate 100 mg/day (N=374) %	Topiramate 200 mg/day (N=501) %	Placebo (N=436) %
Metabolism and Nutrition Disorders				
Anorexia	3.5	7.5	7.2	3.0
Decreased appetite	5.7	7.0	6.8	3.0
Psychiatric Disorders				
Insomnia	4.8	7.0	5.6	3.9
Anxiety	4.0	5.3	5.0	1.8
Expressive language disorder	6.6	5.1	5.2	1.4
Depression	3.5	4.8	7.4	4.1
Depressed mood	0.4	2.9	2.0	0.9
Confusional state	0.4	1.6	2.0	1.1
Mood swings	1.8	1.3	1.0	0.2
Affect lability	0.4	1.1	0.2	0.2
Bradypnea	1.8	1.1	3.4	1.4
Nervous System Disorders				
Paraesthesia	35.7	50.0	48.5	5.0
Dysgeusia	15.4	8.0	12.6	0.9
Hypoaesthesia	5.3	6.7	7.4	1.4
Disturbance in attention	2.6	6.4	9.2	2.3
Somnolence	6.2	5.1	6.8	3.0
Memory impairment	4.0	4.5	6.2	1.6
Amnesia	3.5	2.9	5.2	0.5
Tremor	1.3	1.9	2.4	1.4
Balance disorder	0.4	1.3	0.4	0
Mental impairment	0.4	1.1	1.8	0.9
Eye Disorders				
Vision blurred	4.0	2.4	4.4	2.5
Ear and Labyrinth Disorders				
Tinnitus	0.4	1.3	1.6	0.7
Respiratory, Thoracic and Mediastinal Disorders				
Dyspnoea	1.3	2.7	1.6	1.4
Epistaxis	0.4	1.1	0.6	0.5
Gastrointestinal Disorders				
Nausea	9.3	13.6	14.6	8.3
Diarrhoea	9.3	11.2	10.0	4.4
Dry mouth	1.8	3.2	5.0	2.5
Paraesthesia oral	1.3	2.9	1.6	0.5
Constipation	1.8	2.1	1.8	1.4
Abdominal distension	0	1.3	0.2	0.2
Stomach discomfort	2.2	1.3	1.0	0.2
Gastroesophageal reflux disease	0.4	1.1	1.2	0.5

Table 9: Adverse Drug Reactions Reported by $\geq 1\%$ of Topiramate-Treated Adult Patients in Double-Blind, Placebo-Controlled Migraine Prophylaxis Trials

System/Organ Class Adverse Reaction	Topiramate 50 mg/day (N=227) %	Topiramate 100 mg/day (N=374) %	Topiramate 200 mg/day (N=501) %	Placebo (N=436) %
Musculoskeletal and Connective Tissue Disorders				
Muscle twitching	1.8	1.3	1.8	0.7
General Disorders and Administration Site Conditions				
Fatigue	15.0	15.2	19.2	11.2
Asthenia	0.9	2.1	2.6	0.5
Irritability	3.1	1.9	2.4	0.9
Thirst	1.3	1.6	1.0	0.5
Investigations				
Weight decreased	5.3	9.1	10.8	1.4

Adverse drug reactions reported by $<1\%$ of topiramate-treated adult patients in double-blind, placebo-controlled migraine prophylaxis trials included hypersensitivity, polydipsia, aggression, agitation, anger, crying, disorientation, euphoric mood, hallucination, lack of spontaneous speech, libido decreased, listless, loss of libido, mood altered, panic attack, panic disorder, restlessness, tearfulness, aphasia, burning sensation, clumsiness, cognitive disorder, coordination abnormal, dizziness postural, dyskinesia, dysphasia, hypogeusia, hypokinesia, poor quality sleep, presyncope, psychomotor skills impaired, speech disorder, visual field defect, blepharospasm, diplopia, dry eye, night blindness, visual disturbance, ear discomfort, ear pain, palpitations, flushing, hot flush, dysphonia, nasal congestion, paranasal sinus hypersecretion, epigastric discomfort, gastritis, gingival bleeding, glossodynia, hypoaesthesia oral, alopecia, hypoaesthesia facial, pruritus generalised, urticaria, flank pain, muscular weakness, calculus urinary, micturition urgency, nephrolithiasis, renal pain, sexual dysfunction, feeling abnormal, gait disturbance, malaise, and peripheral coldness.

The recommended dose for migraine prophylaxis is 100 mg/day.

Other Clinical Trial Data

ADRs reported, rate unspecified, in open-label clinical trials of topiramate-treated adult patients are shown in **Table 10**.

Table 10: Adverse Drug Reactions Reported, Rate Unspecified, in Open-Label Clinical Trials of Topiramate-Treated Adult Patients

Nervous System Disorders
Apraxia, aura, complex partial seizure, convulsion, dystonia, grand mal convulsion
Eye Disorders
Glaucoma
Gastrointestinal Disorders
Pancreatitis
General Disorders
Calcinosis

ADRs reported, rate unspecified, in open-label clinical trials of topiramate-treated paediatric patients are shown in **Table 11**.

Table 11: Adverse Drug Reactions Reported, Rate Unspecified, in Open-Label Clinical Trials of Topiramate-Treated Paediatric Patients	
Nervous System Disorders	
	Convulsion, grand mal convulsion
Gastrointestinal Disorders	
	Pancreatitis

Post-marketing Data

Adverse events first identified as ADRs during post-marketing experience with topiramate, presented by frequency category based on spontaneous reporting rates are included in **Table 12**. The frequencies are provided according to the following convention:

Very common: ($\geq 1/10$);

Common: ($\geq 1/100$ to $< 1/10$);

Uncommon: ($\geq 1/1,000$ to $< 1/100$);

Rare: ($\geq 1/10,000$ to $< 1/1000$);

Very rare: ($< 1/10,000$, including isolated reports).

Table 12: Adverse Drug Reactions Identified During Post-marketing Experience with Topiramate by Frequency Category Estimated from Spontaneous Reporting Rates	
Infections and Infestations	
<i>Very rare</i>	Nasopharyngitis
Blood and Lymphatic System Disorders	
<i>Very rare</i>	Neutropenia
Immune System Disorders	
<i>Very rare</i>	Allergic oedema
Metabolism and Nutrition Disorders	
<i>Very rare</i>	Hyperammonemia
<i>Very rare</i>	Hyperammonemic encephalopathy
Psychiatric Disorders	
<i>Very rare</i>	Feeling of despair
Eye Disorders	
<i>Very rare</i>	Abnormal sensation in eye
<i>Very rare</i>	Angle closure glaucoma
<i>Very rare</i>	Conjunctival oedema
<i>Very rare</i>	Eye movement disorder
<i>Very rare</i>	Eyelid oedema
<i>Very rare</i>	Maculopathy
<i>Very rare</i>	Myopia
Respiratory, Thoracic and mediastinal Disorders	
<i>Very rare</i>	Cough
Skin and Subcutaneous Tissue Disorders	
<i>Very rare</i>	Erythema multiforme

Table 12: Adverse Drug Reactions Identified During Post-marketing Experience with Topiramate by Frequency Category Estimated from Spontaneous Reporting Rates	
Infections and Infestations	
<i>Very rare</i>	Periorbital oedema
<i>Very rare</i>	Stevens-Johnson syndrome
<i>Very rare</i>	Toxic epidermal necrolysis
Musculoskeletal and Connective Tissue Disorders	
<i>Very rare</i>	Joint swelling
<i>Very rare</i>	Limb discomfort
Renal and Urinary Disorders	
<i>Very rare</i>	Renal tubular acidosis
<i>Very rare</i>	Nephrocalcinosis
General Disorders and Administration Site Reactions	
<i>Very rare</i>	Generalised oedema
<i>Very rare</i>	Influenza like illness
Investigations	
<i>Very rare</i>	Weight increased

Reporting Suspected Adverse Effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at www.tga.gov.au/reporting-problems.

4.9 OVERDOSE

Signs and Symptoms

Ingestion of between 6 and 40 g topiramate have been reported in a few patients. Signs and symptoms included: headache, agitation, drowsiness, lethargy, convulsions, speech disturbances, blurred vision, diplopia, mentation impaired, abnormal coordination, stupor, hypotension, abdominal pain, dizziness, depression and hypokalaemia. The clinical consequences were not severe in most cases, but deaths have been reported after polydrug overdoses involving topiramate.

Topiramate overdose can result in severe metabolic acidosis (see **SECTION 4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE - METABOLIC ACIDOSIS**).

A patient who ingested a dose calculated to be between 96 and 110 g topiramate was admitted to hospital with coma lasting 20-24 hours followed by full recovery after 3 to 4 days.

Treatment

General supportive measures are indicated and an attempt should be made to remove undigested drug from the gastro intestinal tract using activated charcoal.

Treatment should be appropriately supportive. Haemodialysis has been shown to be an effective means of removing topiramate from the body. The patient should be well hydrated.

For information on the management of overdose, contact the Poisons Information Centre on 13 11 26 (Australia).

5 PHARMACOLOGICAL PROPERTIES

5.1 PHARMACODYNAMIC PROPERTIES

Mechanism of Action

Topiramate is classified as a sulfamate substituted monosaccharide. Three pharmacological properties of topiramate have been identified that may contribute to its anticonvulsant activity:

- Topiramate reduces the frequency at which action potentials are generated when neurons are subjected to a sustained depolarisation, which is indicative of a state dependent blockage of voltage sensitive sodium channels.
- Topiramate markedly enhances the activity of GABA at some types of GABA_A receptors. This effect was not blocked by flumazenil, a benzodiazepine antagonist, nor did topiramate increase the duration of the channel open time, differentiating topiramate from barbiturates that modulate GABA_A receptors. Because the antiepileptic profile of topiramate differs markedly from that of the benzodiazepines, it may modulate a benzodiazepine insensitive subtype of GABA_A receptor.
- Topiramate antagonises the ability of kainate to activate the kainate/AMPA subtype of excitatory amino acid (glutamate) receptors but has no apparent effect on the activity of N-methyl-D-aspartate (NMDA) at the NMDA receptor subtype.

In addition, topiramate inhibits some isoenzymes of carbonic anhydrase. This pharmacological effect is much weaker than that of acetazolamide, a known carbonic anhydrase inhibitor, and is not thought to be a major component of topiramate's antiepileptic activity.

The mechanism(s) of action of topiramate in migraine prophylaxis is unknown.

Clinical Trials

Monotherapy Epilepsy

Three double blind, randomised, parallel group clinical trials were conducted to evaluate the efficacy and safety of topiramate given as monotherapy. Study YI and EPMN-104 evaluated the safety and efficacy of topiramate monotherapy using a dose-response design by comparing the low dose regimen with the high dose regimen. Study EPMN-105 compared topiramate monotherapy to carbamazepine or valproate in patients with newly diagnosed epilepsy.

In study YI, adults with refractory partial onset seizures (n = 48) were converted from their existing treatment to TAMATE 100 mg/day or 1,000 mg/day as monotherapy. The high dose group was statistically superior to the low dose group for efficacy variables. 54% of high dose patients achieved monotherapy compared with 17% in the low dose group with the difference between the doses being statistically significant (p = 0.005). The mean time to exit was significantly greater in the high dose group (p = 0.002). The investigator and subject global evaluations of clinical response statistically favoured the high dose group (less than or equal to 0.002).

In study EPMN-104, adult and paediatric patients with recently diagnosed epilepsy (n = 252) were randomised into the low dose (25 or 50 mg/day) or the high dose group (200 or 500 mg/day) based on their bodyweight. Overall, 54% of high dose patients and 39% of low dose patients were reported to be seizure free during the double blind phase (p = 0.022). The high dose group was also superior to the low dose group with respect to seizure frequency distribution (p = 0.008) and the difference in time to first seizure across three plasma topiramate concentration strata (p = 0.015).

In study EPMN-105, patients with newly diagnosed epilepsy (n = 613) were randomised to receive either 100 or 200 mg/day of topiramate or standard epileptic treatment (carbamazepine or valproate).

Topiramate was at least as efficacious as carbamazepine or valproate in reducing seizures in these patients; the 95% confidence intervals for the difference between the two treatment groups were narrow and included zero, indicating that there was no statistically significant difference between both groups. The two treatment groups were also comparable with respect to all clinical utility and efficacy endpoints including time to exit, proportion of seizure free subjects and time to first seizure.

Patients (n = 207; 32 were aged less than or equal to 16 years) who completed the double blind phase of study YI and EMPN-104 were enrolled in long-term extension studies with the majority of patients receiving topiramate for two to five years. In these studies, sustained efficacy was demonstrated with long-term administration of topiramate as monotherapy. There was no significant change in dosage during the extension period and no indication that effectiveness of topiramate monotherapy diminished with continued exposure.

The safety profile of topiramate in monotherapy trials is consistent with that of the add-on trials.

Add-on Therapy Epilepsy

Over 2,000 patients worldwide were involved in the clinical trials of topiramate as an add-on treatment in adults and children with the following type of epilepsy: partial onset seizures with or without secondary generalised seizures, primary generalised tonic-clonic seizures and seizures associated with Lennox-Gastaut syndrome. These trials were randomised, placebo controlled, double blind, multicentre, parallel group studies in which patients were given topiramate or placebo as add-on treatment while they were receiving phenytoin, carbamazepine, primidone, phenobarbitone or valproic acid, as concomitant therapy.

These trials had 4 to 12 weeks as the 'run in' phases, several weeks of titration and then up to 12 weeks of stabilisation. Topiramate reduced monthly seizure rates and increased responder rates (fraction of patients with at least 50% seizure reduction) significantly compared to placebo. In addition, Topiramate significantly reduced seizure severity in patients with Lennox-Gastaut syndrome. No evidence of tolerance to Topiramate has been demonstrated in humans.

In a pooled analysis of two clinical trials involving patients with primary generalised tonic-clonic epilepsy, topiramate (n = 79) was statistically better than placebo (n = 81) (p = 0.004). In these two trials, 17 patients who were 16 years or younger received topiramate.

There is limited clinical experience with topiramate at or above daily dose of 1000 mg. Comparative data or data on the safety and efficacy of using topiramate with lamotrigine, gabapentin or vigabatrin are not available. Elderly patients and patients with known or suspected coronary artery disease did not participate in these studies.

Migraine

The clinical development program to evaluate the efficacy of topiramate in the prophylaxis of migraine included four double blind, placebo controlled, parallel group trials. Each trial started with a washout period (14 to 28 days) for subjects already taking prophylactic drugs, followed by a 28 day 'run in' phase, an eight week dose titration phase and a 12 or 18 week maintenance phase.

The pooled results of the two pivotal trials, evaluating topiramate doses of 50 (n = 233), 100 (n = 244) and 200 mg/day (n = 288), found a median percent reduction in average monthly migraine period rate of 35, 51 and 49% respectively, compared to 21% for the pooled placebo group (n = 229). Notably 27% of patients administered topiramate 100 mg/day achieved at least a 75% reduction in migraine frequency, while 52% achieved at least a 50% reduction.

Study MIGR-003 demonstrated that topiramate 100 mg/day was comparable in terms of efficacy to propranolol 160 mg/day. There was no statistically significant difference between the two groups in the primary efficacy endpoint or clinically significant 50% responder rate (43% for propranolol 160 mg/day, 37% for topiramate 100 mg/day (-6% difference, 95% CI (-17%, +6%), p = 0.28), 35% for topiramate 200 mg/day (-7% difference, 95% CI (-19%, +4%), p = 0.17).

Results from each trial are summarised in **Table 13**.

Table 13: Responder Rates (at least a 50% reduction in average monthly migraine period compared to baseline –ITT)				
Study	Placebo	50mg/day	100mg/day	200mg/day
MIGR-001	23%	36% p≤0.05* 12% ^a (1%, 24%) ^b	54% p≤0.001* 31% ^a (19%, 42%) ^b	52% p≤0.001* 29% ^a (17%, 41%) ^b
MIGR-002	23%	39% p≤0.05* 16% ^a (4%, 28%) ^b	49% p≤0.001* 26% ^a (15%, 38%) ^b	47% p≤0.001* 24% ^a (12%, 36%) ^b
MIGR-003	22%		37% p≤0.05* 15% ^a (4%, 25%) ^b	35% p≤0.05* 13% ^a (2%, 23%) ^b
CAPSS-155	34%			40% NS 6% ^a (-8%, 19%) ^b

*Nominal p values for comparison of topiramate with placebo.
^adifference – treatment responder rate of topiramate minus placebo.
^b95% Confidence interval – pairwise difference of topiramate minus placebo.
 NOTE: The overall safety profile of topiramate observed in the migraine studies was generally consistent with that established for epilepsy therapy.

5.2 PHARMACOKINETIC PROPERTIES

The pharmacokinetic profile of topiramate compared to other antiepileptic drugs shows a long plasma elimination half-life, linear pharmacokinetics, predominantly renal clearance, absence of significant protein binding and lack of clinically relevant active metabolites. Topiramate is not a potent inducer of drug metabolising enzymes. Routine monitoring of plasma topiramate concentrations is not necessary and it can be administered without regard to meals. In clinical studies, there was no consistent relationship between plasma concentrations and efficacy or adverse events.

Topiramate was rapidly and well absorbed and distributed in total body water following oral administration in animals. The same metabolic and elimination pathways were present as in human subjects. C_{max} values were similar to those obtained in human subjects but topiramate was more rapidly cleared in animals resulting in lower overall systemic exposure.

Absorption

Based on the recovery of radioactivity from urine in humans, the mean extent of absorption of a 100 mg dose of ¹⁴C-topiramate was at least 81%. Following oral administration of 100 mg topiramate to healthy subjects, a mean peak plasma concentration (C_{max}) of approximately 1.5 microgram/mL was achieved with two or three hours (T_{max}). The bioavailability of topiramate is not significantly affected by food.

In a separate study involving 24 normal, healthy, non-smoking male and female subjects under fasting conditions, the mean absorption rate of a 200 mg dose of topiramate was found to be approximately 132.65 microgram/mL. Following oral administration of 200 mg topiramate to healthy subjects, a mean peak plasma concentration (C_{max}) of approximately 4.5 microgram/mL was achieved within two hours of administration.

Distribution

Generally 13 to 17% of topiramate is bound to plasma proteins. A low capacity binding site for topiramate in/on erythrocytes that is saturated at steady state has been observed. Following single dose administration, the volume of distribution varies inversely with dose. The mean apparent volume of distribution has been

measured as 0.55 to 0.8 L/kg for a single dose of 100 to 1,200 mg. There is an effect of gender on the volume of distribution. Values for females are about 50% lower than those of males. This is attributed to the higher percent body fat in females and is of no clinical consequence.

Metabolism

Topiramate is not extensively metabolised (approximately 20%) in healthy volunteers. It is metabolised up to 50% in patients receiving concomitant antiepileptic therapy with known inducers of drug metabolising enzymes. Six metabolites formed through hydroxylation, hydrolysis and glucuronidation have been isolated, characterised and identified in plasma, urine and faeces of humans. Each metabolite represents less than 3% of the total radioactivity excreted following administration of ¹⁴C-topiramate. Two metabolites, which retained most of the structure of topiramate, were tested and found to have little to no anticonvulsant activity.

Excretion

In humans, the major route of elimination of unchanged topiramate and its metabolites is via the kidney (at least 81% of the dose). Approximately 66% of a dose of ¹⁴C-topiramate was excreted unchanged in the urine within four days. Following twice a day dosing with 50 and 100 mg of topiramate, the mean renal clearance was approximately 18 and 17 mL/minute respectively. There is evidence of renal tubular reabsorption of topiramate. This is supported by studies in rats where topiramate was co-administered with probenecid and a significant increase in renal clearance of topiramate was observed. Overall, plasma clearance is approximately 20 to 30 mL/minute in humans following oral administration. Concomitant multiple dose administration of topiramate, 100 to 400 mg twice a day, with phenytoin and carbamazepine shows dose proportional increases in plasma concentrations of topiramate.

Topiramate exhibits low intersubject variability in plasma concentrations and therefore, has predictable pharmacokinetics. The pharmacokinetics of topiramate are linear at steady state with plasma clearance remaining constant and area under the plasma concentration curve (AUC) increasing in a dose proportional manner over a 200 to 800 mg daily oral dose range. Patients with normal renal function may take four to eight days to reach steady-state plasma concentrations. The main C_{max} following multiple, twice a day oral doses of 100 mg to healthy subjects was 6.76 microgram/mL. Following administration of multiple doses of 50 and 100 mg of topiramate twice a day, the mean plasma elimination half-life was approximately 21 hours.

Following administration of a single dose of 200 mg of topiramate in 24 subjects under fasting conditions, the mean plasma elimination half-life was approximately 28 hours.

Patients with renal impairment

The plasma and renal clearance of topiramate decreased in patients with moderate and severe impaired renal function (creatinine clearance < 70 mL/minute). As a result, higher steady-state plasma concentrations are expected for a given dose in renally impaired patients compared to those with normal renal function. In addition, patients with renal impairment will require a longer time to reach steady-state at each dose (see **section 4.2 DOSE AND METHOD OF ADMINISTRATION**).

Topiramate is effectively removed from plasma by haemodialysis. A prolonged period of haemodialysis may cause topiramate concentration to fall below levels that are required to maintain an anti-seizure effect. To avoid rapid drops in topiramate plasma concentration during haemodialysis, a supplemental dose of topiramate may be required. The actual adjustment should take into account 1) the duration of dialysis period, 2) the clearance rate of the dialysis system being used, and 3) the effective renal clearance of topiramate in the patient being dialysed.

Patients with hepatic impairment

Plasma clearance of topiramate decreased a mean of 26% in patients with moderate to severe hepatic impairment. Therefore, topiramate should be administered with caution in patients with hepatic impairment.

Elderly

Plasma clearance of topiramate is unchanged in elderly subjects in the absence of underlying renal disease or hepatic impairment. Patients more than 71 years of age have not been studied.

Paediatric up to 12 years

The pharmacokinetics of topiramate are linear in children receiving the drug as add-on therapy. The clearance is independent of dose and steady-state plasma concentrations increase proportionally to dose. Hepatic enzyme inducing antiepileptic drugs decrease the steady-state plasma concentrations. In comparison to adults, however, children have a higher clearance and shorter elimination half-life when topiramate is used as adjunctive therapy to both enzyme inducing and non-enzyme inducing antiepileptic drugs. Consequently, the plasma concentrations of topiramate for the same mg/kg dose may be lower in children compared to adults.

5.3 PRECLINICAL SAFETY DATA

Genotoxicity

Topiramate was not genotoxic in a series of assays for gene mutations, chromosomal damage or DNA damage.

Carcinogenicity

No evidence of carcinogenicity was seen in rats following oral administration of topiramate for 2 years at doses of 120mg/kg. An increased incidence of urinary bladder tumours of a proliferative nature was observed in mice following oral administration of topiramate for 22 months at doses of 300mg/kg. These tumours probably resulted from chronic irritation and may lack clinical significance. The plasma concentration exposure obtained in the animal studies was less than the likely clinical exposure at the maximum recommended dose.

6 PHARMACEUTICAL PARTICULARS

6.1 LIST OF EXCIPIENTS

The tablets also contain the following inactive excipients: microcrystalline cellulose, povidone, colloidal anhydrous silica, sodium starch glycollate type A, and magnesium stearate.

The coating contained in the tablets are as follows:

25 mg: OPADRY complete film coating system YS-1-7003 White (PI 11956)

50 mg: OPADRY complete film coating system 03B92164 Yellow (PI 12654)

100 mg: OPADRY complete film coating system 03B92180 Yellow (PI 12655)

200 mg: OPADRY complete film coating system 05B16131 Maroon (PI 12656)

6.2 INCOMPATIBILITIES

Incompatibilities were either not assessed or not identified as part of the registration of this medicine.

6.3 SHELF LIFE

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

6.4 SPECIAL PRECAUTIONS FOR STORAGE

TAMATE tablets should be stored in a dry place below 25°C, and protected from light and moisture.

6.5 NATURE AND CONTENTS OF CONTAINER

Container type: blister pack Al/Al

Pack sizes: 60's

6.6 SPECIAL PRECAUTIONS FOR DISPOSAL


In Australia, any unused medicine or waste material should be disposed of by taking it to your local pharmacy.

6.7 PHYSICOCHEMICAL PROPERTIES

Topiramate is a white crystalline powder with a bitter taste.

Topiramate is most soluble in alkaline solutions with a pH of 9 to 10, including those that contain sodium hydroxide or sodium phosphate. It is freely soluble in acetone, chloroform, dimethylsulfoxide and ethanol. The solubility in water is 9.8 mg/mL. Its saturated solution has a pH of 6.3.

Chemical Structure:



Chemical Formula: 2,3:4,5-bis-0-(1-methylethylidene)-β-D-fructopyranose sulfamate

Molecular formula: C₂₁H₂₁NO₈S

Molecular weight: 339.36

CAS Number: (CAS) Registry Number:97240-79-4

7 MEDICINE SCHEDULE (POISONS STANDARD)

S4 (Prescription Only Medicine)

8 SPONSOR

Alphapharm Pty Limited

Level 1, 30 The Bond

30 – 34 Hickson Road

Millers Point NSW 2000

ABN 93 002 359 739

www.mylan.com.au

9 DATE OF FIRST APPROVAL

Date of first inclusion in the Australian Register of Therapeutic Goods: 01/06/2009

10 DATE OF REVISION

Date of the most recent TGA approved changes: 09/04/2019

Summary Table of Changes

Section Changed	Summary of New Information
All sections	Reformatted in line with the revised Australian form for providing product information format
4.4	- Addition of a warning for women of childbearing potential
4.5	- Addition of cross references and information for metabolic acidosis (risk of nephrolithiasis or nephrocalcinosis)
4.6	- Addition of an interaction with vitamin K-antagonist anticoagulant medications
4.8	- Additional information regarding the dose dependent nature of topiramate with the risk of congenital malformations and SGA; Addition of pre-term labor and premature delivery; Addition of effects observed in breastfed infants. Addition of ‘nephrocalcinosis’

TAMATE_pi\04/19/04