

# OPSUMIT®

(OP-suh-mit)

Macitentan (ma-SEE-ten-tan) 10 mg Tablets

## Consumer Medicine Information

---

### What is in this leaflet

This leaflet answers some common questions about OPSUMIT.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using this medicine against the benefits they expect it will have for you.

**If you have any concerns about using this medicine, speak to your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

---

### What OPSUMIT is used for

OPSUMIT tablets contain macitentan which belongs to the class of medicines called "endothelin receptor antagonists".

OPSUMIT is used for the treatment of pulmonary arterial hypertension (PAH), it can be used on its own or with other drugs to treat PAH. PAH is high blood pressure in the blood vessels (the pulmonary arteries) that carry blood from the heart to the lungs. In people with PAH, these arteries get narrower, so the heart has to work harder to pump blood through them. This causes people to feel tired, dizzy, and short of breath.

OPSUMIT widens the pulmonary arteries, making it easier for the heart to pump blood through them. This

lowers the blood pressure and relieves the symptoms and improves the course of the disease.

Your doctor however, may prescribe OPSUMIT for another purpose.

**Ask your doctor if you have any questions about why it has been prescribed for you.**

---

### Before you take OPSUMIT

#### **When you must not take OPSUMIT**

Do NOT take OPSUMIT if you are pregnant or if you are planning to become pregnant, or if you could become pregnant because you are not using reliable birth control (contraception). Talk to your doctor about what birth control methods are reliable whilst taking OPSUMIT.

Tell your doctor if you are breast-feeding. You are advised to stop breast-feeding if OPSUMIT is prescribed for you, because it is not known whether this medicine passes into human breast milk.

Male fertility - the potential risk for humans is unknown. Talk to your doctor if you have any questions or concerns about this.

**Do not take OPSUMIT if you have ever had an allergic reaction to macitentan or any of the ingredients listed at the end of this leaflet.**

Symptoms of an allergic reaction may be mild or severe. They usually include some or all of the following:

wheezing, swelling of the lips/mouth, difficulty in breathing, hay fever, lumpy rash (hives) or fainting.

**OPSUMIT contains small amounts of a sugar called lactose. If you have an intolerance to lactose or any other sugars tell your doctor.**

**Do not take OPSUMIT if you have liver disease or if you have very high levels of liver enzymes in your blood. Talk to your doctor, who will decide if this medicine is suitable for you.**

**Do not use OPSUMIT after the expiry date [EXP.] printed on the pack.**

If you use it after the expiry date has passed, it may not work as well.

**Do not use OPSUMIT if the packaging is torn or shows signs of tampering.**

If you're not sure whether you should be using OPSUMIT, talk to your doctor. The safety and efficacy of OPSUMIT has not been proven in patients under 12 years of age.

#### **Before you start to take OPSUMIT**

**You must tell your doctor if:**

- **you are allergic to foods, dyes, preservatives or any other medicines,**
- **you are pregnant, trying or become pregnant,**
- **you are breast-feeding,**
- **you have liver problems,**
- **you have kidney problems,**
- **you have low blood pressure,**
- **you have anaemia or low red blood cells.**

## **Blood Tests**

Your doctor will also order some blood tests before you start treatment with OPSUMIT and also during treatment with OPSUMIT, this is to test:

- whether you have anaemia (a reduced number of red blood cells)
- whether your liver is working properly.

Signs that your liver may not be working properly include:

- nausea (urge to vomit)
- vomiting
- fever (high temperature)
- pain in your stomach (abdomen)
- jaundice (yellowing of your skin or the whites of your eyes)
- dark-coloured urine
- itching of your skin
- lethargy or fatigue (unusual tiredness or exhaustion)
- flu-like syndrome (joint and muscle pain with fever)

**If you notice any of these signs, tell your doctor immediately.**

## **Pregnancy and Breast-feeding**

Don't take OPSUMIT if you or your partner are pregnant or planning to become pregnant as it may harm unborn babies conceived before, during or soon after treatment.

You must not become pregnant for at least 3 months after stopping OPSUMIT.

If you are a woman who could become pregnant, your doctor will ask you to take a pregnancy test before you take OPSUMIT.

If you are a male you should avoid exposing your partner to your semen by use of appropriate contraception.

If it's possible you could become pregnant, use at least two reliable forms of birth control (contraception) while you're taking OPSUMIT. You must continue to use contraception

for at least 3 months after stopping OPSUMIT. Talk to your doctor about this.

If you do become pregnant, talk to your doctor immediately.

Don't breast feed while you are taking OPSUMIT.

## **Taking Other medicines**

**You must tell your doctor if you are taking:**

- rifampicin, clarithromycin, (used to treat infections)
- phenytoin (a medicine used to treat seizures)
- carbamazepine (a medicine used to treat depression and epilepsy)
- St John's Wort (a herbal preparation used to treat depression)
- ritonavir, saquinavir (used to treat HIV infections)
- ketoconazole (except shampoo), itraconazole, voriconazole (medicines used to treat fungal infections)

**You are taking any other medicines, including medicines you buy without a prescription from a pharmacy, supermarket or health food shop.**

Some medicines may affect the way other medicines work. Your doctor or pharmacist will be able to tell you what to do when using OPSUMIT with other medicines.

---

## **How to take OPSUMIT**

---

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

### **How much to take**

The recommended dose of OPSUMIT for adults and children over 12 years of age and weighing more than 40 kg is one 10 mg tablet, once a day. Swallow the whole

tablet, with a glass of water, do not chew or break the tablet.

You can take OPSUMIT with or without food.

### **How long to take it**

Use OPSUMIT for as long as your doctor advises you to. OPSUMIT is generally used over a prolonged period of time, possibly years. It should not be stopped suddenly.

### **If you use too much (overdose)**

If you have taken more tablets than you have been told to take, ask your doctor for advice.

### **If you forget to take it**

If you forget to take OPSUMIT, take a dose as soon as you remember, then continue to take your tablets at the usual times. Do not take a double dose to make up for a forgotten tablet.

---

## **While you are using OPSUMIT**

---

### **Things you must do**

**Tell your doctor or pharmacist that you are taking OPSUMIT if you are about to start on any new medicines.**

**Tell your doctor if you become pregnant or are trying to become pregnant.**

**Tell your doctor if, for any reason, you have not used your medicine exactly as prescribed.**

### **Things you must not do**

OPSUMIT is a treatment that you will need to keep on taking to control your PAH. Do not stop taking OPSUMIT unless you have agreed this with your doctor.

If you have any further questions on the use of OPSUMIT, ask your doctor or pharmacist.

**Do not give this medicine to anyone else, even if their symptoms seem similar to yours.**

**Do not use OPSUMIT to treat any other complaints unless your doctor says to.**

### **Things to be careful of**

As with many other medicines, OPSUMIT may cause headaches in some people.

**Be careful driving or operating machinery until you know how OPSUMIT affects you. If you are affected, do not drive or operate machinery.**

---

## **Side Effects**

---

**Check with your doctor as soon as possible if you have any problems while taking OPSUMIT, even if you do not think the problems are connected with the medicine or are not listed in this leaflet.**

Like all medicines, OPSUMIT can cause side effects in some people.

**Tell your doctor at once if you experience any of the following while you are receiving OPSUMIT:**

### **Very common side effects**

- anaemia (low number of red blood cells) or haemoglobin decreases
- headache
- bronchitis (inflammation of the airways)
- nasopharyngitis (inflammation of the throat & nasal passages)
- oedema/fluid retention (swelling)

### **Common side effects**

- diarrhoea
- abdominal pain
- pharyngitis (inflammation of the throat)
- influenza (flu)
- urinary tract infection (bladder infection)

- hypotension (low blood pressure)
- nasal congestion
- pruritus (severe itching of the skin)

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

**Do not be alarmed by this list of possible side effects.**

**You may not experience any of them.**

This is not a complete list of all possible side-effects. Others may occur in some people and there may be some side-effects not yet known.

**Tell your doctor if you notice anything else that is making you feel unwell, even if it is not on this list.**

**Ask your doctor if you don't understand anything in this list.**

---

## **Storing OPSUMIT**

---

### **Storage**

**Keep this medicine where young children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

**Do not leave them in the car or on window sills.**

**Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.**

Do not store above 30°C.

Medicines should not be disposed of in wastewater or household waste.

Ask your pharmacist how to dispose of medicines you no longer require. These measures will help to protect the environment.

---

## **Product description**

---

### **What it looks like**

OPSUMIT 10 mg tablets are white to off white, biconvex, round, film-coated tablets with "10" on two sides.

### **Pack size**

OPSUMIT is supplied as 10 mg film-coated tablets in blister packs of, 9 or 30 tablets.

### **Ingredients**

#### **Active ingredient:**

Each tablet of OPSUMIT contains 10 mg of macitentan.

#### **Inactive ingredients:**

- lactose 38.86mg
- microcrystalline cellulose
- povidone
- sodium starch glycollate type A
- magnesium stearate
- polysorbate 80
- polyvinyl alcohol
- titanium dioxide
- purified talc
- lecithin
- xanthan gum.

### **Supplier**

OPSUMIT® is supplied in Australia by Actelion Pharmaceuticals Australia Pty Limited.

Suite 6

13b Narabang Way

Belrose NSW 2085

OPSUMIT is distributed in NZ by Pharmacy Retailing (NZ) Ltd t/a Healthcare Logistics

58 Richard Pearse Drive

Airport Oaks, Auckland NZ.

This leaflet was prepared in February 2018.

OPSUMIT 10 mg tablets packs:

- blister packs of 9 tablets AUST  
R 205624
- blister packs of 30 tablets  
AUST R 205624