

TRELEGY ELLIPTA

100/62.5/25

Fluticasone furoate 100 micrograms, umeclidinium (as bromide) 62.5 micrograms and vilanterol (as trifenate) 25 micrograms per inhalation

Consumer Medicine Information

What is in this leaflet

Please read this leaflet carefully before you start using TRELEGY ELLIPTA.

This leaflet answers some common questions about TRELEGY ELLIPTA.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking TRELEGY ELLIPTA against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What TRELEGY ELLIPTA is used for

TRELEGY ELLIPTA is used to treat chronic obstructive pulmonary disease (COPD). TRELEGY ELLIPTA is a dry powder inhaler. When you breathe in using the inhaler, the medicine is deposited into the lungs.

COPD is a long-term condition that slowly gets worse. Symptoms include shortness of breath, cough, chest discomfort and coughing up mucus.

TRELEGY ELLIPTA contains three active ingredients: fluticasone furoate, umeclidinium (as bromide) and vilanterol (as trifenate).

Fluticasone furoate belongs to a group of medicines called corticosteroids, often simply called steroids. They are not 'anabolic steroids' which are the steroids sometimes misused by athletes.

Corticosteroids are used to reduce inflammation. They reduce the swelling and irritation in the small air passages in the lungs and so ease breathing problems.

Umeclidinium and vilanterol belong to a group of medicines called bronchodilators. They work together to help open the airways and make it easier for air to get in and out of the lungs.

When TRELEGY ELLIPTA is used regularly, it can help to control the breathing difficulties related to your disease and help with the effects of the disease on your everyday life.

TRELEGY ELLIPTA should not be used to relieve a sudden attack of breathlessness or wheezing. If you get this sort of attack you must use a quick-acting inhaler (such as VENTOLIN).

Your doctor may have prescribed TRELEGY ELLIPTA for another reason.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

This medicine is not addictive.

This medicine is available only with a doctor's prescription.

TRELEGY ELLIPTA should not be used in children or adolescents under the age of 18 years.

Before you use TRELEGY ELLIPTA

When you must not use it

Don't use TRELEGY ELLIPTA:

- **if you are allergic (hypersensitive) to lactose or milk protein**
- **if you are allergic (hypersensitive) to fluticasone furoate, umeclidinium, vilanterol or any other ingredients of TRELEGY ELLIPTA (listed at the end of this leaflet).**

If you think either of these applies to you, don't use TRELEGY ELLIPTA until you have checked with your doctor.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing, coughing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching, redness or hives on the skin
- suddenly feeling weak or light headed (may lead to collapse or loss of consciousness).

TRELEGY ELLIPTA contains lactose monohydrate.

If you have been diagnosed with an intolerance to some sugars, or to milk protein, talk to your doctor before you use TRELEGY ELLIPTA.

TRELEGY ELLIPTA is not usually recommended for use during pregnancy.

If you are pregnant, if you think you may be pregnant or if you are planning to have a baby, don't use TRELEGY ELLIPTA without asking your doctor.

Your doctor will consider the benefit to you and the risk to your baby of taking TRELEGY ELLIPTA while you are pregnant.

If you are breast-feeding, check with your doctor before you take TRELEGY ELLIPTA.

It is not known whether the ingredients of TRELEGY ELLIPTA can pass into breast milk.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to use it

Tell your doctor if you have or have had any of the following medical conditions:

- asthma
- heart problems or high blood pressure
- liver disease, as you may be more likely to have side effects
- being or have been treated for tuberculosis (TB) or pneumonia
- eye problems such as glaucoma or cataracts
- an enlarged prostate, difficulty passing urine or a blockage in your bladder

- weak bones (osteoporosis)
- any type of viral, bacterial or fungal infection
- any other medical conditions
- a problem with your immune system.

Check with your doctor before you use TRELEGY ELLIPTA if you think any of these apply to you.

Your doctor can discuss with you the risks and benefits involved.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop. This should include all of the medicines that you are using for your COPD.

Some medicines and TRELEGY ELLIPTA may interfere with each other. These include:

- medicines called beta blockers, to treat high blood pressure or other heart problems
- ketoconazole, to treat fungal infections
- ritonavir, to treat HIV
- other long acting medicines similar to this medicine that are used to treat breathing problems (i.e. inhalers used as maintenance or preventer therapy)
- medicines to treat depression or mood/mental disorders (such as monoamine oxidase inhibitors or tricyclic antidepressants).

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to use TRELEGY ELLIPTA

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions in the user leaflet, ask your doctor or pharmacist for help.

How much to use

Always use TRELEGY ELLIPTA exactly as your doctor has told you. Check with your doctor, nurse or pharmacist if you're not sure.

The dose of TRELEGY ELLIPTA is one inhalation once daily at the same time each day.

Don't use more than your doctor tells you to use.


How to use the inhaler

The full instructions for using TRELEGY ELLIPTA are given on a leaflet inside the pack. A brief summary of the instructions is provided below.

TRELEGY ELLIPTA is ready to use straight away. No preparation or checks of the inhaler are required.

Step 1: Prepare a dose

- Wait to open the cover until you are ready to take your dose. Slide the cover fully down until you hear a "click".



Do not open TRELEGY ELLIPTA until you are ready to take a dose.

Step 2: Inhale your medication

- Whilst holding the inhaler away from your mouth, breathe out as far as is comfortable.
- Put the mouthpiece between your lips, and close your lips firmly around it.




- Take one long, steady, deep breath in. Hold this breath for about 3-4 seconds or as long as is comfortable.
- Remove the inhaler from your mouth.
- Breathe out slowly and gently away from the mouthpiece.

After using TRELEGY ELLIPTA, you may clean the mouthpiece, using a dry tissue, before you close the cover. Do not immerse TRELEGY ELLIPTA in water.

Step 3: Close the inhaler and rinse your mouth

- Slide the cover upwards as far as it will go, to cover the mouthpiece.



When to use it

Use TRELEGY ELLIPTA regularly.

It is very important that you use TRELEGY ELLIPTA every day, as instructed by your doctor. This will help to keep you free of symptoms throughout the day and night.

If you feel you are getting breathless or wheezy more often than normal, or if you are using a quick-acting inhaler (such as VENTOLIN) more than usual, see your doctor.

How long to use it

Use TRELEGY ELLIPTA for as long as your doctor recommends. It will only be effective as long as you are using it. Don't stop unless your doctor advises you to, even if you feel better.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and use your next dose when you are meant to. Otherwise, use it as soon as you remember, then go back to using it as you would normally.

Don't take an extra dose to make up for a missed dose.

If you are not sure what to do, ask your doctor or pharmacist.

If you become wheezy or breathless, use your quick-acting inhaler (e.g. VENTOLIN), then seek medical advice.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

In Australia, immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, if you think that you or anyone else may have taken too much TRELEGY ELLIPTA. Do this even if there are no signs of discomfort or poisoning.

If you have used larger doses than instructed for a long period of time, it is particularly important that you ask your doctor or pharmacist for advice. This is because larger doses of TRELEGY ELLIPTA may reduce the amount of steroid hormones produced naturally by your body.

While you are using TRELEGY ELLIPTA

Things you must do

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking TRELEGY ELLIPTA.

Tell any other doctors, dentists, and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.

If you become pregnant while taking this medicine, tell your doctor immediately.

Keep all of your doctor's appointments so that your progress can be checked.

Things you must not do

Do not take TRELEGY ELLIPTA to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your medicine or lower the dosage without checking with your doctor.

Check with your doctor if you are unsure about which medicines for COPD you should continue taking whilst being treated with TRELEGY ELLIPTA. You should stop taking any long-acting inhalers which contain inhaled corticosteroids or bronchodilators. However, you can continue to use a quick-acting inhaler (such as VENTOLIN) during an acute episode.

Things to be careful of

Be careful driving or operating machinery until you know how TRELEGY ELLIPTA affects you.

Side effects

Like all medicines, TRELEGY ELLIPTA can cause side effects, although not everybody gets them.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Common side effects

These may affect up to 1 in 10 people:

- common cold
- headache
- cough
- joint pain
- back pain
- itchy, runny or blocked nose
- flu (influenza)
- infection of the nose, sinuses or throat
- infection of the upper airways
- infection of the lungs (pneumonia)*

*affected approximately 1 in 50 people in clinical trials. In general, risk factors for pneumonia in people with COPD receiving inhaled medicines containing a corticosteroid include smoking, prior pneumonia, low body weight and severe COPD.

Uncommon side effects

These may affect up to 1 in 100 people:

- sore, raised patches in the mouth or throat caused by a fungal infection (candidiasis). Rinsing your mouth out with water immediately after using TRELEGY ELLIPTA may help stop this side effect developing
- irregular heart beat
- faster heart beat
- pain in the back of the mouth and throat

- weakening of the bones, leading to fractures.

If you think you are having an allergic reaction to TRELEGY ELLIPTA, stop using this medicine and tell your doctor immediately or go the accident and emergency department at your nearest hospital. Symptoms of an allergic reaction usually include some or all of the following:

- shortness of breath
- wheezing, coughing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching, redness or hives on the skin
- suddenly feeling weak or light headed (may lead to collapse or loss of consciousness).

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

After using TRELEGY ELLIPTA

Storage

Do not use TRELEGY ELLIPTA after the expiry date shown on the pack.

Store in the original package container in order to protect from moisture and do not open the foil lid until ready to inhale for the first time.

Safely throw away TRELEGY ELLIPTA one month after you open the foil tray or when the counter reads "0", whichever comes first. Write the date the inhaler should be discarded on the label in the space provided. The date should be added as soon as the inhaler has been removed from the tray.

Keep your inhaler in a cool dry place where the temperature stays below 30°C.

If you store in a refrigerator allow the inhaler to return to room temperature for at least an hour before use.

Do not store TRELEGY ELLIPTA or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

What it looks like

TRELEGY ELLIPTA is inhaled through the mouth using the ELLIPTA device. The active ingredients are in separate blisters in powder form inside the device. There are either 14 or 30 blisters on each strip, and so each device contains either 14 or 30 doses depending on which pack size you have been given.

The ELLIPTA device itself is a plastic inhaler with a light grey body, a beige mouthpiece cover and a dose counter. It is packaged in a foil laminate tray with a peelable lid foil. The tray contains a desiccant sachet, to reduce moisture in the packaging. Once you have opened the lid of the tray, throw the desiccant away - do not open, eat or inhale it.

Ingredients

The active ingredients in TRELEGY ELLIPTA are fluticasone furoate, umeclidinium (as bromide) and vilanterol (as trifenate).

Each dose contains 100 micrograms of fluticasone furoate, 62.5 micrograms of umeclidinium (as bromide) and 25 micrograms of vilanterol (as trifenate).

TRELEGY ELLIPTA also contains the inactive ingredients:

- lactose monohydrate
- magnesium stearate.

Supplier

TRELEGY ELLIPTA is supplied in Australia by:

GlaxoSmithKline Australia Pty Ltd
Level 4, 436 Johnston Street,
Abbotsford, Victoria 3067,
Australia.

Where to go for further information

Pharmaceutical companies are not in a position to give people an individual diagnosis or medical advice. Your doctor or pharmacist is the best person to give you advice on the treatment of your condition. You may also be able to find general information about your disease and its treatment from patient information groups and books, for example in public libraries.

Information on COPD is available from the website of the Lung Foundation Australia (<https://lungfoundation.com.au/>).

A copy of this CMI is available from the GSK Australia website (www.gsk.com.au/trelegy).

The information provided applies only to TRELEGY ELLIPTA.

TRELEGY ELLIPTA: AUST R
284636

Trade marks are owned by or licensed to the GSK group of companies.

© 2017 GSK group of companies or licensor.

This leaflet was prepared on 16 January 2018.

Version 1.0