

## PRODUCT INFORMATION


**KALYDECO®**

(ivacaftor)

### NAME OF THE MEDICINE

Ivacaftor is *N*-(2,4-di-tert-butyl-5-hydroxyphenyl)-1,4-dihydro-4-oxoquinoline-3-carboxamide. The empirical formula of ivacaftor is C<sub>24</sub>H<sub>28</sub>N<sub>2</sub>O<sub>3</sub> and its molecular weight is 392.49. The pKa values of ivacaftor are 9.40 and 11.60. The log D value of ivacaftor is 5.68 at pH=7.4 and 25°C.

Its structural formula is:



CAS Number: 873054-44-5

### DESCRIPTION

Ivacaftor is a white to off-white powder that is practically insoluble in water (<0.05 microgram/mL).

#### *Tablets*

KALYDECO is available as a light blue, capsule-shaped, film-coated tablet for oral administration containing 150 mg of ivacaftor per tablet. Each tablet contains the following inactive ingredients: microcrystalline cellulose, lactose monohydrate, hypromellose acetate succinate, croscarmellose sodium, sodium lauryl sulfate, silicon dioxide, magnesium stearate, carnauba wax, Opadry II complete film coating system 85F90614 Blue (PI# 108371), OPACODE monogramming ink S-1-17823 BLACK (PI# 12108).

#### *Granules*

KALYDECO is available as white to off-white granules enclosed in unit dose sachet for oral administration containing 50 mg or 75 mg of ivacaftor per sachet. Each sachet contains the following inactive ingredients: lactose monohydrate, hypromellose acetate succinate, croscarmellose sodium, sodium lauryl sulfate, silicon dioxide, magnesium stearate, mannitol, sucralose.

### PHARMACOLOGY

#### **Mechanism of Action**

Ivacaftor is a selective potentiator of the Cystic Fibrosis Transmembrane conductance Regulator (CFTR) protein. *In vitro*, ivacaftor increases the open probability of the CFTR channel gate to enhance chloride transport. This has been demonstrated in normal CFTR and in mutant forms of CFTR that have reduced channel-open probability, such as G551D-CFTR and R117H-CFTR. However, the exact mechanism leading ivacaftor to prolong the gating activity of some mutant CFTR forms has not been completely elucidated.

### **Pharmacodynamics**

In clinical trials (Studies 1 and 2) in patients with the *G551D* mutation in one allele of the *CFTR* gene administered 150 mg of ivacaftor tablets every 12 hours, Study 1 in patients 12 years and older and Study 2 in patients 6 to 11 years of age, ivacaftor led to rapid (15 days), substantial (the mean change in sweat chloride from baseline through Week 24 was -48 mmol/L [95% CI -51, -45] and -54 mmol/L [95% CI -62, -47] respectively), and sustained (through 48 weeks) reduction in sweat chloride concentration.

In a clinical trial in patients aged 6 years and older administered 150 mg of ivacaftor tablets every 12 hours who had a non-*G551D* gating mutation in the *CFTR* gene (Study 5, Part 1), treatment with ivacaftor led to a rapid (15 days) and substantial mean change in sweat chloride from baseline of -49 mmol/L (95% CI -57, -41) after 8 weeks of treatment.

In a clinical trial (Study 6) in 69 patients age 6 years or older with CF who had an *R117H* mutation in the *CFTR* gene, the treatment difference in mean change in sweat chloride from baseline through 24 weeks of treatment was -24 mmol/L (95% CI -28, -20) and in patients 18 years or older it was -22 mmol/L (95% CI -26, -17). The mean sweat chloride change was consistent across subgroups, including age, poly-T status, and FEV<sub>1</sub>.

In an open-label, Phase 3 clinical study in 34 patients aged 2 to less than 6 years administered either 50 mg or 75 mg of ivacaftor granules every 12 hours (Study 7), the mean absolute change from baseline in sweat chloride was -47 mmol/L (95% CI -58, -36) at Week 24.

### **Pharmacokinetics**

The pharmacokinetics of ivacaftor are similar between healthy adult volunteers and patients with CF. After oral administration of a single 150-mg dose to healthy volunteers in a fed state, the mean ( $\pm$ SD) for AUC and C<sub>max</sub> were 10600 (5260) ng\*hr/mL and 768 (233) ng/mL, respectively. The apparent terminal half-life was approximately 12 hours following a single dose in the fed state. The apparent clearance (CL/F) of ivacaftor was similar for healthy subjects and patients with CF. The mean ( $\pm$ SD) of CL/F for the 150 mg dose was 17.3 (8.4) L/hr in healthy subjects at steady state.

After every 12-hour dosing, steady-state plasma concentrations of ivacaftor were reached by Days 3 to 5, with an accumulation ratio ranging from 2.2 to 2.9.

### **Absorption**

Following multiple oral dose administrations of ivacaftor, the exposure of ivacaftor generally increased with a dose from 25 mg every 12 hours to 450 mg every 12 hours. The exposure of ivacaftor increased approximately 2.5- to 4-fold when given with fat-containing food. Therefore, ivacaftor should be administered with fat-containing food. The median (range) t<sub>max</sub> is approximately 4.0 (3.0, 6.0) hours in the fed state.

Ivacaftor granules (2 x 75-mg sachets) had similar bioavailability to the 150-mg ivacaftor tablet when given with fat-containing food to adult patients. The effect of food on ivacaftor absorption is similar for ivacaftor granules and the 150-mg tablet formulation.

### **Distribution**

Ivacaftor is approximately 99% bound to plasma proteins, primarily to alpha 1-acid glycoprotein and albumin. Ivacaftor does not bind to human red blood cells.

After oral administration of 150 mg every 12 hours for 7 days in healthy volunteers in a fed state, the mean ( $\pm$ SD) apparent volume of distribution was 353 (122) L.

### **Metabolism**

Ivacaftor is extensively metabolised in humans. *In vitro* and *in vivo* data indicate that ivacaftor is primarily metabolised by CYP3A. M1 and M6 are the two major metabolites of ivacaftor in humans. M1 has approximately one-sixth the potency of ivacaftor and is considered pharmacologically active. M6 has less than one-fiftieth the potency of ivacaftor and is not considered pharmacologically active.

### **Excretion**

Following oral administration, the majority of ivacaftor (88%) was excreted in the faeces after metabolic conversion. The major metabolites M1 and M6 accounted for approximately 65% of the total dose excreted with 22% as M1 and 43% as M6. There was negligible urinary excretion of ivacaftor as unchanged parent and minimal urinary excretion (6.6%) of ivacaftor plus metabolites.

### **Special Populations**

#### *Children and Adolescents*

Predicted ivacaftor exposure based on observed ivacaftor concentrations in Phase 2 and 3 studies as determined using population pharmacokinetic (PK) analysis is presented by age group (and body weight for patients less than 12 years of age) in Table 1. Exposures in 6- to 11-year-olds are predictions based on simulations from the population PK model using data obtained for this age group.

<b>Age Group</b>	<b>Dose</b>	<b>C<sub>min, ss</sub> (ng/mL) (Median)</b>	<b>C<sub>min, ss</sub> (ng/mL) (Mean [SD])</b>	<b>AUC<sub>ss</sub> (ng*h/mL) (Median)</b>	<b>AUC<sub>ss</sub> (ng*h/mL) (Mean [SD])</b>
2- to 5-year-olds (<14 kg)	50 mg q12h	536	577 (317)	9840	10500 (4260)
2- to 5-year-olds (≥14 kg to <25 kg)	75 mg q12h	580	629 (296)	10200	11300 (3820)
6- to 11-year-olds (≥14 kg to <25 kg)	75 mg q12h	584	641 (329)	10050	10760 (4470)
6- to 11-year-olds (≥25 kg)	150 mg q12h	838	958 (546)	13920	15300 (7340)
12- to 17-year-olds	150 mg q12h	508	564 (242)	8670	9240 (3420)
Adults (≥18 years old)	150 mg q12h	634	701 (317)	9840	10700 (4100)

#### *Gender*

The effect of gender on ivacaftor pharmacokinetics was evaluated using population pharmacokinetics of data from clinical studies of ivacaftor. No dose adjustments are necessary based on gender.

#### *Renal Impairment*

Pharmacokinetic studies have not been performed with ivacaftor in patients with renal impairment. In a human pharmacokinetic study, there was minimal elimination of ivacaftor and its metabolites in urine (only 6.6% of total radioactivity was recovered in the urine). There was negligible urinary excretion of ivacaftor as unchanged parent (less than 0.01% following a single oral dose of 500 mg). Therefore, no dose adjustments are recommended for mild and moderate renal impairment. However, caution is recommended when administering ivacaftor to patients with severe renal impairment (creatinine clearance less than or equal to

30 mL/min) or end-stage renal disease (see DOSAGE AND ADMINISTRATION and PRECAUTIONS).

#### *Hepatic Impairment*

Following a single dose of 150 mg of ivacaftor, adult patients with moderately impaired hepatic function (Child-Pugh Class B, score 7 to 9) had similar ivacaftor  $C_{max}$  (mean [ $\pm$ SD] of 735 [331] ng/mL), but an approximately 2-fold increase in ivacaftor  $AUC_{0-\infty}$  (mean [ $\pm$ SD] of 16800 [6140] ng\*hr/mL) compared with healthy subjects matched for demographics. Simulations for predicting the steady-state exposure of ivacaftor showed that by reducing the dosage from 150 mg q12h to 150 mg once daily, adults with moderate hepatic impairment would have comparable steady-state  $C_{min}$  values as those obtained with a dose of 150 mg q12h in adults without hepatic impairment. Therefore, a reduced dose of one tablet or one sachet once daily is recommended in patients with moderate hepatic impairment. The impact of mild hepatic impairment (Child-Pugh Class A, score 5 to 6) on the pharmacokinetics of ivacaftor has not been studied, but the increase in ivacaftor  $AUC_{0-\infty}$  is expected to be less than two-fold. Therefore, no dose adjustment is necessary for patients with mild hepatic impairment.

Studies have not been conducted in patients with severe hepatic impairment (Child-Pugh Class C, score 10 to 15), but exposure is expected to be higher than in patients with moderate hepatic impairment. The use of KALYDECO in patients with severe hepatic impairment is therefore not recommended unless the benefits outweigh the risks. In such cases, the starting dose should be one tablet or one sachet every other day. Dosing intervals should be modified according to clinical response and tolerability (see DOSAGE AND ADMINISTRATION and PRECAUTIONS).

## **CLINICAL TRIALS**

The efficacy of KALYDECO has been evaluated in six clinical trials including two Phase 3, randomised, double-blind, placebo-controlled, multi-centre studies of clinically stable patients with CF who had the *G551D* mutation in the *CFTR* gene on at least 1 allele and had FEV<sub>1</sub> (forced expiratory volume exhaled in the first second)  $\geq$ 40% predicted.

Patients in both studies were randomised 1:1 to receive either 150 mg of KALYDECO tablets or placebo every 12 hours with fat-containing food for 48 weeks in addition to their prescribed CF therapies (e.g., tobramycin, dornase alfa). The use of inhaled hypertonic saline was not permitted.

### **Study 1: Study in Patients with CF ( $\geq$ 12 years) with a *G551D-CFTR* Mutation**

Study 1 evaluated 161 patients who were 12 years of age or older; 122 (75.8%) patients had the *F508del* mutation in the second allele. At the start of the study, patients in the placebo group used some medicinal products at a higher frequency than the ivacaftor group. These medications included dornase alfa (73.1% versus 65.1%), salbutamol (53.8% versus 42.2%), tobramycin (44.9% versus 33.7%), and salmeterol/fluticasone (41.0% versus 27.7%). At baseline, mean predicted FEV<sub>1</sub> was 63.6% (range: 31.6% to 98.2%) and mean age was 26 years (range: 12 to 53 years).


### **Study 2: Study in Patients with CF (6 to 11 years) with a *G551D-CFTR* Mutation**

Study 2 evaluated 52 patients who were 6 to 11 years of age at screening; mean (SD) body weight was 30.9 (8.63) kg; 42 (80.8%) of patients had the *F508del* mutation in the second allele. At baseline, mean predicted FEV<sub>1</sub> was 84.2% (range: 44.0% to 133.8%) and mean age was 9 years (range: 6 to 11 years); 8 (30.8%) patients in the placebo group and 4 (15.4%) patients in the ivacaftor group had an FEV<sub>1</sub> less than 70% predicted at baseline.

The primary efficacy endpoint in both studies was the mean absolute change from baseline in percent predicted FEV<sub>1</sub> through 24 weeks of treatment.

The treatment difference between ivacaftor and placebo for the mean absolute change (95% CI) in percent predicted FEV<sub>1</sub> from baseline through Week 24 was 10.6 percentage points (8.6, 12.6) in Study 1 and 12.5 percentage points (6.6, 18.3) in Study 2 (Figure 1). The treatment difference between ivacaftor and placebo for the mean relative change (95% CI) in percent predicted FEV<sub>1</sub> from baseline through Week 24 was 17.1% (13.9, 20.2) in Study 1 and 15.8% (8.4, 23.2) in Study 2. The mean change from baseline through Week 24 in FEV<sub>1</sub> (L) was 0.37 L in the ivacaftor group and 0.01 L in the placebo group in Study 1 and 0.30 L in the ivacaftor group and 0.07 L in the placebo group in Study 2. In both studies, improvements in percent predicted FEV<sub>1</sub> were rapid in onset (Day 15  $P < 0.0001$  and  $P = 0.0004$  for Study 1 and 2, respectively) and durable through 48 weeks ( $P < 0.0001$  and  $P = 0.0017$  for Study 1 and 2, respectively).

**Figure 1: Mean Absolute Change from Baseline in Percent Predicted FEV<sub>1</sub>**



FEV<sub>1</sub> results at 24 and 48 weeks by age subgroups are shown in Table 2.

Subgroup Age (years)	Study	Through Week 24 (percentage points) (95% CI)	Through Week 48 (percentage points) (95% CI)
6 to 11	2	12.5 (6.6, 18.3)	10.0 (4.5, 15.5)
12 to 17	1	11.9 (5.9, 17.9)	11.4 (5.4, 17.4)
≥18	1	9.9 (7.8, 12.0)	9.9 (7.7, 12.0)

CI: confidence interval; FEV<sub>1</sub>: forced expiratory volume in 1 second; MMRM: mixed-effects model for repeated measures

The results for clinically relevant secondary endpoints are shown in Table 3.

<b>Table 3. Effect of Ivacaftor on Other Efficacy Endpoints in Studies 1 and 2</b>				
<b>Endpoint</b>	<b>Study 1</b>		<b>Study 2</b>	
	<b>Treatment difference<sup>a</sup> (95% CI)</b>	<b>P value</b>	<b>Treatment difference<sup>a</sup> (95% CI)</b>	<b>P value</b>
<b>Mean absolute change from baseline in CFQ-R<sup>b</sup> respiratory domain score (points)<sup>c</sup></b>				
Through Week 24	8.1 (4.7, 11.4)	<0.0001	6.1 (-1.4, 13.5)	0.1092
Through Week 48	8.6 (5.3, 11.9)	<0.0001	5.1 (-1.6, 11.8)	0.1354
<b>Relative risk of pulmonary exacerbation</b>				
Through Week 24	0.40 <sup>d</sup>	0.0016	NA	NA
Through Week 48	0.46 <sup>d</sup>	0.0012	NA	NA
<b>Mean absolute change from baseline in body weight (kg)</b>				
At Week 24	2.8 (1.8, 3.7)	<0.0001	1.9 (0.9, 2.9)	0.0004
At Week 48	2.7 (1.3, 4.1)	0.0001	2.8 (1.3, 4.2)	0.0002
<b>Mean absolute change from baseline in BMI (kg/m<sup>2</sup>)</b>				
At Week 24	0.94 (0.62, 1.26)	<0.0001	0.81 (0.34, 1.28)	0.0008
At Week 48	0.93 (0.48, 1.38)	<0.0001	1.09 (0.51, 1.67)	0.0003
<b>Mean change from baseline in z-scores</b>				
Weight-for-age z-score at Week 48 <sup>e</sup>	0.33 (0.04, 0.62)	0.0260	0.39 (0.24, 0.53)	<0.0001
BMI-for-age z-score at Week 48 <sup>e</sup>	0.33 (0.002, 0.65)	0.0490	0.45 (0.26, 0.65)	<0.0001
CI: confidence interval; NA: not analysed due to low incidence of events				
<sup>a</sup> Treatment difference = effect of ivacaftor – effect of placebo				
<sup>b</sup> CFQ-R: Cystic Fibrosis Questionnaire-Revised is a disease-specific, health-related quality-of-life measure for CF. Minimum Clinically Important Difference (MCID) = 4 units.				
<sup>c</sup> Study 1 data were pooled from CFQ-R for adults/adolescents and CFQ-R for children 12 to 13 years of age; Study 2 data were obtained from CFQ-R for children 6 to 11 years of age.				
<sup>d</sup> Hazard ratio for time to first pulmonary exacerbation				
<sup>e</sup> In patients under 20 years of age (CDC growth charts)				

### **Study 3: Study in Patients (≥12 years) with CF with the *F508del* Mutation in the *CFTR* Gene**

Study 3 (Part A) was a 16-week, 4:1 randomised, double-blind, placebo-controlled, parallel-group, Phase 2 study of ivacaftor (150 mg tablets every 12 hours) in 140 patients with CF age 12 years and older who were homozygous for the *F508del* mutation in the *CFTR* gene and who had FEV<sub>1</sub> ≥40% predicted (see PRECAUTIONS).

The mean absolute change from baseline through Week 16 in percent predicted FEV<sub>1</sub> (primary efficacy endpoint) was 1.5 percentage points in the ivacaftor group and -0.2 percentage points in the placebo group. The estimated treatment difference for ivacaftor versus placebo was 1.7 percentage points (95% CI -0.6, 4.1); this difference was not statistically significant (*P*=0.15).


#### Study 4: Open-Label Extension Study in Patients ≥6 years

Study 4 is an open-label extension study to evaluate the safety and efficacy of long-term treatment of orally administered ivacaftor (150 mg tablets every 12 hours) in patients continuing from Studies 1 and 2. It enrolled 144 adolescents/adults who completed Study 1 (age ≥12 years) and 48 children who completed Study 2 (age 6 to 11 years). The percent predicted FEV<sub>1</sub> range at the beginning of Study 4 was 29.1% to 126.7%. The use of inhaled hypertonic saline was permitted. A pre-specified analysis of all patients' data was performed after all patients from Studies 1 and 2 who had not discontinued received 96 weeks of treatment with ivacaftor in Study 4 (for a total exposure of 144 weeks for patients who received ivacaftor in Studies 1 and 2).


In Study 4, the substantial lung function improvement and weight gain (as seen in Studies 1 and 2) (Table 4) and CFQ-R changes (as seen in Study 1) persisted through 144 weeks of cumulative KALYDECO treatment. The time-to-first pulmonary exacerbation showed a consistent trend across 144 weeks for patients treated with KALYDECO from Study 1. Too few patients experienced pulmonary exacerbations in Study 2 to perform a meaningful statistical analysis.

<b>Table 4. Effect of Ivacaftor in Study 4</b>					
<b>Original study and treatment group</b>	<b>Duration of KALYDECO treatment (weeks)</b>	<b>Absolute change in percent predicted FEV<sub>1</sub> (percentage points)</b>		<b>Absolute change in body weight (kg)</b>	
		<b>N</b>	<b>Mean (SD)</b>	<b>N</b>	<b>Mean (SD)</b>
<b>Study 1</b>					
<b>KALYDECO</b>	<b>48*</b>	<b>77</b>	<b>9.4 (8.3)</b>	<b>77</b>	<b>3.4 (4.9)</b>
	<b>144</b>	<b>72</b>	<b>9.4 (10.8)</b>	<b>72</b>	<b>4.1 (7.1)</b>
<b>Placebo</b>	<b>0*</b>	<b>67</b>	<b>-1.2 (7.8)<sup>†</sup></b>	<b>67</b>	<b>0.3 (2.7)<sup>†</sup></b>
	<b>96</b>	<b>55</b>	<b>9.5 (11.2)</b>	<b>55</b>	<b>3.0 (4.7)</b>
<b>Study 2</b>					
<b>KALYDECO</b>	<b>48*</b>	<b>26</b>	<b>10.2 (15.7)</b>	<b>26</b>	<b>6.1 (2.9)</b>
	<b>144</b>	<b>25</b>	<b>10.3 (12.4)</b>	<b>25</b>	<b>14.8 (5.7)</b>
<b>Placebo</b>	<b>0*</b>	<b>22</b>	<b>-0.6 (10.1)<sup>†</sup></b>	<b>22</b>	<b>2.9 (1.8)<sup>†</sup></b>
	<b>96</b>	<b>21</b>	<b>10.5 (11.5)</b>	<b>21</b>	<b>10.1 (4.1)</b>
* Treatment occurred during blinded, controlled, 48-week, Phase 3 study.					
† Change from prior study baseline after 48 weeks of placebo treatment.					


**Figure 2: Mean Absolute Change in Percent Predicted FEV<sub>1</sub> from Baseline in Studies 1 and 2 to Week 144 in Study 4**



**Figure 3: Absolute Change in Weight in Studies 1 and 2 to Week 144 in Study 4**



**Figure 4: Absolute Change in Body Mass Index (BMI) in Studies 1 and 2 to Week 144 in Study 4**



**Study 5: Study in Patients (≥6 years) with CF with non-G551D Gating Mutations**

Study 5 was a Phase 3, two-part, randomised, double-blind, placebo-controlled, crossover study (Part 1) with an open-label extension period (Part 2) to evaluate the efficacy and safety of ivacaftor in patients with CF who have a non-G551D gating mutation in the *CFTR* gene (*G178R*, *S549N*, *S549R*, *G551S*, *G970R*, *G1244E*, *S1251N*, *S1255P*, or *G1349D*). Patients who completed Part 1 of this study (randomised, double-blind, placebo-controlled, 8-week crossover) continued into the 16-week open-label Part 2 of the study.

Study 5 evaluated 39 patients with CF who were 6 years of age or older (mean age 23 years) with baseline FEV<sub>1</sub> ≥40% predicted (mean FEV<sub>1</sub> 78% predicted [range: 43% to 119%]).

In Part 1, patients were randomised 1:1 to receive either 150 mg of ivacaftor tablets or placebo every 12 hours with fat-containing food for 8 weeks in addition to their prescribed CF therapies during the first Treatment Period and crossed over to the other treatment for the second 8 weeks. The two 8-week Treatment Periods were separated by a 4- to 8-week Washout Period. The use of inhaled hypertonic saline was not permitted.

The primary efficacy endpoint was improvement in lung function as determined by the mean absolute change from baseline in percent predicted FEV<sub>1</sub> through 8 weeks of treatment.

In Part 1 of Study 5, the observed treatment difference between ivacaftor and placebo for the mean absolute change in percent predicted FEV<sub>1</sub> from baseline through Week 8 was 10.7 percentage points (*P*<0.0001). Improvements in percent predicted FEV<sub>1</sub> were observed regardless of age, disease severity, sex, geographic region, and *Pseudomonas aeruginosa* infection status at baseline. Improvement in percent predicted FEV<sub>1</sub> was rapid in onset (Day 15, *P*<0.0001) and durable through 8 weeks (*P*<0.0001) of treatment with ivacaftor.

Treatment with ivacaftor resulted in substantial, consistent, and statistically significant treatment effects across the secondary endpoints of the absolute change from baseline in BMI and BMI-for-age z-score at Week 8 was 0.7 kg/m<sup>2</sup> (*P*<0.0001) and 0.3 points (*P*=0.0010), respectively, and CFQ-R respiratory domain score through Week 8 was 9.6 points (*P*=0.0004; Minimum Clinically Important Difference = 4 units) when compared to placebo. Together, these results demonstrate the positive effects of ivacaftor treatment on pulmonary and extrapulmonary measures.

### **Study 6: Study in Patients with CF with an *R117H* Mutation in the *CFTR* Gene**

The efficacy and safety of KALYDECO in patients with CF who have an *R117H* mutation in the *CFTR* gene were evaluated in a randomized, double-blind, placebo-controlled, parallel-group clinical trial of patients aged 6 years and older with cystic fibrosis. Fifty-nine of 69 patients completed 24 weeks of treatment. Two patients discontinued and 8 patients did not complete treatment due to study termination. Patients who were 12 years and older had FEV<sub>1</sub> at screening between 40-90% predicted, and patients who were 6-11 years of age had FEV<sub>1</sub> at screening between 40-105% predicted. The patients had well preserved BMIs (mean overall: 23.76 kg/m<sup>2</sup>) and a high proportion were pancreatic sufficient as assessed by a low rate of pancreatic enzyme replacement therapy use (pancreatin: 11.6%; pancrelipase: 5.8%). Patients who had persistent *Burkholderia cenocepacia*, *Burkholderia dolosa*, or *Mycobacterium abscessus* isolated from sputum at screening, and those with abnormal liver function defined as 3 or more liver function tests (ALT, AST, AP, GGT, total bilirubin)  $\geq 3$  times the ULN, were excluded.

Patients were randomized 1:1 to receive either 150 mg of KALYDECO (n=34) or placebo (n=35) every 12 hours with food containing fat for 24 weeks in addition to their prescribed CF therapies.

The primary efficacy endpoint was improvement in lung function as determined by the mean absolute change from baseline in percent predicted FEV<sub>1</sub> through 24 weeks of treatment. The treatment difference for absolute change in percent predicted FEV<sub>1</sub> through Week 24 was 2.1 percentage points (analysis conducted with the full analysis set which included all 69 patients), and did not reach statistical significance.

In a subgroup analysis in patients 18 years and older, the treatment difference for the mean absolute change from baseline through Week 24 in percent predicted FEV<sub>1</sub> was 5.0 percent points (95% CI 1.1, 8.8). In a subgroup analysis in patients 6-11 years of age, the treatment difference for the mean absolute change from baseline through Week 24 in percent predicted FEV<sub>1</sub> was -6.3 percentage points (95% CI -12.0, -0.7). No statistical analysis was conducted for subjects 12 to 17 years of age because only 2 patients were enrolled in the clinical study (Table 4).

Other efficacy variables that were analyzed included absolute change in sweat chloride from baseline through Week 24, improvement in cystic fibrosis respiratory symptoms through Week 24 as assessed by the CFQ-R respiratory domain score (Table 4), absolute change in body mass index (BMI) at Week 24, and time to first pulmonary exacerbation. The overall treatment difference for the absolute change from baseline in BMI at Week 24 was 0.3 kg/m<sup>2</sup> and the calculated hazard ratio for time to first pulmonary exacerbation was 0.93, which were not statistically significant.

Statistically significant improvements in clinical efficacy (FEV<sub>1</sub>, CFQ-R respiratory domain) were seen in several subgroup analyses, and decreases in sweat chloride were observed in all subgroups. Subgroups analyzed included those based on age, lung function, and poly-T status (Table 5).

<b>Table 5. Effect of KALYDECO on Overall Population (Percent Predicted FEV<sub>1</sub>, CFQ-R Respiratory Domain Score, and Sweat Chloride) and in Relevant Subgroups Through 24 Weeks</b>										
<b>Absolute Change Through Week 24* - All Randomized Patients</b>										
		<b>% Predicted FEV<sub>1</sub> (Percentage Points)</b>			<b>CFQ-R Respiratory Domain Score (Points)</b>			<b>Sweat Chloride (mmol/L)</b>		
<b>Subgroup Parameter</b>	<b>Study Drug</b>	<b>n</b>	<b>Mean</b>	<b>Treatment Difference (95% CI)</b>	<b>n</b>	<b>Mean</b>	<b>Treatment Difference (95% CI)</b>	<b>n</b>	<b>Mean</b>	<b>Treatment Difference (95% CI)</b>
<b><i>R117H – All Patients</i></b>										
	Placebo	35	0.5	2.1	34	-0.8	8.4	35	-2.3	-24.0
	Kalydeco	34	2.6	(-1.1, 5.4)	33	7.6	(2.2, 14.6)	32	-26.3	(-28.0, -19.9)
<b><i>Subgroup by Age</i></b>										
<b>6-11</b>	Placebo	8	3.5	-6.3	7	-1.6	-6.1	8	1.0	-27.6
	Kalydeco	9	-2.8	(-12.0, -0.7)	8	-7.7	(-15.7, 3.4)	8	-26.6	(-37.2, -18.1)
<b>12-17</b>	Placebo	1	---	---	1	---	---	1	---	---
	Kalydeco	1	---	---	1	---	---	1	---	---
<b>≥18</b>	Placebo	26	-0.5	5.0	26	-0.5	12.6	26	-4.0	-21.9
	Kalydeco	24	4.5	(1.1, 8.8)	24	12.2	(5.0, 20.3)	23	-25.9	(-26.5, -17.3)
<b><i>Subgroup by Poly-T Status<sup>†</sup></i></b>										
<b>5T</b>	Placebo	24	0.7	5.3	24	-0.6	15.3	24	-4.6	-24.2
	Kalydeco	14	6.0	(1.3, 9.3)	14	14.7	(7.7, 23.0)	13	-28.7	(-30.2, -18.2)
<b>7T</b>	Placebo	5	-0.9	0.2	5	-6.0	5.2	5	3.9	-24.1
	Kalydeco	11	-0.7	(-8.1, 8.5)	11	-0.7	(-13.0, 23.4)	10	-20.2	(-33.9, -14.3)
<b><i>Subgroup by Baseline FEV<sub>1</sub> % Predicted</i></b>										
<b>&lt;70%</b>	Placebo	15	0.4	4.0	15	3.0	11.4	15	-3.8	-25.5
	Kalydeco	13	4.5	(-2.1, 10.1)	13	14.4	(1.2, 21.6)	12	-29.3	(-31.8, -19.3)
<b>70-90%</b>	Placebo	14	0.2	2.6	13	-3.6	8.8	14	-3.1	-20.0
	Kalydeco	14	2.8	(-2.3, 7.5)	14	5.2	(-2.6, 20.2)	14	-23.0	(-26.9, -12.9)
<b>&gt;90%</b>	Placebo	6	2.2	-4.3	6	-2.5	-0.7	6	1.0	-26.8
	Kalydeco	7	-2.1	(-9.9, 1.3)	6	-3.2	(-10.4, 9.0)	6	-25.9	(-39.5, -14.1)
* MMRM analysis with fixed effects for treatment, age, week, baseline value, treatment by week, and subject as a random effect										
† (n=54) Poly-T status confirmed by genotyping										

The efficacy of ivacaftor beyond 2 years of treatment has not been examined in clinical trials in children with *R117H* mutation.

### **Study 7: Study in Paediatric Patients with CF aged 2 to less than 6 years with *G551D* or another gating mutation**

The pharmacokinetic profile, safety and efficacy of Ivacaftor granules in patients aged 2 through less than 6 years with CF who have a *G551D* mutation or other gating (class III) mutation in the *CFTR* gene were studied in a 2-part, 24-week, open-label, Phase 3 clinical study in 34 patients (efficacy was a tertiary endpoint). The results of studies 1, 2 and 5 which examined Ivacaftor in older subjects were used as supportive data.

Patients in Study 7 were 2 to less than 6 years of age (mean age 3 years). The majority of the study population had pancreatic insufficiency; 25 of the 27 patients with available data had Fecal Elastase-1 values <50 µg/g at baseline. The mean Fecal Elastase-1 value at baseline was 28 µg/g. Patients with evidence of colonization with *Burkholderia cenocepacia*, *Burkholderia dolosa*, or *Mycobacterium abscessus* and those with abnormal liver function defined as 3 or more liver function tests (ALT, AST, AP, GGT, total bilirubin) ≥3 x ULN at screening were excluded.

Patients were assigned to the ivacaftor 50-mg or ivacaftor 75-mg groups according to their weight at enrolment (patients weighing less than 14 kg received ivacaftor 50-mg granules and patients weighing more than 14 kg received ivacaftor 75-mg granules). Ivacaftor was administered orally every 12 hours with fat-containing food in addition to their prescribed CF therapies.

Efficacy endpoints evaluated were absolute change from baseline in sweat chloride through 24 weeks of treatment (see Pharmacodynamic effects), measures of nutritional status including absolute change from baseline in weight, body mass index (BMI), and stature (supported by weight, BMI, and stature-for-age z-scores) at 24 weeks of treatment, and measures of pancreatic function including Fecal Elastase-1 and Immunoreactive Trypsinogen (IRT) results at 24 weeks.

Overall, sweat chloride (see Pharmacodynamic effects), and mean absolute changes from baseline in nutritional status (weight, BMI, stature; supported by weight, BMI, and stature-for-age z-scores), showed improvement from baseline in Study 7. Changes in weight and BMI z-scores are consistent with results shown in 6- to 11-year-olds in Study 2 and Study 5. Substantial increases from baseline in Fecal Elastase-1 were observed at Week 24. The mean (SD) overall change from baseline in Faecal Elastase-1 (n=27) was 99.8 µg/g (138.4). Six patients with initial levels below 200µg/g achieved, at week 24, a level of ≥200 µg/g. Changes from baseline in IRT (mean baseline value 33.58 ng/mL) showed decreases at Week 24. Results are shown in Table 6. Data on percent predicted FEV<sub>1</sub> (exploratory endpoint) were available for 3 patients in the ivacaftor 50 mg group and 17 patients in the 75 mg dosing group. The mean (SD) overall change in percent predicted FEV<sub>1</sub> from baseline at week 24 (exploratory endpoint) was 1.8 (17.81).

<b>Table 6: Effect of Ivacaftor on Efficacy Endpoints in Study 7</b>			
<b>Endpoint</b>	<b>Week 24 Mean (SD) Absolute Changes From Baseline</b>		
	<b>Ivacaftor 50 mg (N=10)</b>	<b>Ivacaftor 75 mg (N=24)</b>	<b>Overall (N=34)</b>
<b>Secondary Endpoints</b>			
Weight (kg)	1.0 (0.42)	1.5 (0.55)	1.4 (0.56)
BMI (kg/m <sup>2</sup> )	0.33 (0.54)	0.31 (0.55)	0.32 (0.54)
Stature (cm)	2.5 (1.45)	3.5 (0.93)	3.3 (1.17)
<b>Other Endpoints</b>			
Weight-for-age z-scores (unit)	0.18 (0.32)	0.21 (0.23)	0.20 (0.25)
BMI-for-age z-scores (unit)	0.46 (0.46)	0.34 (0.42)	0.37 (0.42)
Stature-for-age z-scores (unit)	-0.25 (0.45)	0.08 (0.22)	-0.01 (0.33)
Fecal Elastase-1 (µg/g)*	128 (191.84)	93.5 (128.28)	99.8 (138.35)
Immunoreactive Trypsinogen (IRT)†	-24.37 (21.71)	-19.54 (25.11)	-20.70 (23.99)
* N=5, ivacaftor 50 mg; N=22, ivacaftor 75 mg; N=27, overall			
† N=7, ivacaftor 50 mg; N=21 ivacaftor 75 mg; N=28, overall			

## INDICATIONS

KALYDECO is indicated for the treatment of cystic fibrosis (CF) in patients aged 2 years and older who have a *G551D* or other gating (class III) mutation in the *CFTR* gene.

KALYDECO is indicated for the treatment of cystic fibrosis (CF) in patients aged 6 years and older who have an *R117H* mutation in the *CFTR* gene.

## **CONTRAINDICATIONS**

In cases of hypersensitivity to the active substance or to any component of this medication, patients should not be treated with this medicine.

## **PRECAUTIONS**

Efficacy results from a Phase 2 study in patients with CF who were homozygous for the *F508del* mutation in the *CFTR* gene showed no statistically significant difference in FEV<sub>1</sub> over 16 weeks of ivacaftor treatment compared to placebo (see CLINICAL TRIALS). Therefore, use of KALYDECO in these patients is not recommended.

### **Effects on Fertility**

Ivacaftor impaired fertility and reproductive performance indices in male and female rats at an oral dose of 200 mg/kg/day (yielding approximately 10 and 5 times, respectively, the systemic exposure anticipated in patients at the maximum recommended human dose (MRHD) based on summed AUCs of ivacaftor and its major metabolites) when dams were dosed prior to and during early pregnancy. The pregnancy rate was decreased, oestrus cycling was disrupted and pre-implantation loss was increased. These effects occurred in the presence of significant maternal toxicity. No effects on male or female fertility and reproductive performance indices were observed at  $\leq 100$  mg/kg/day (yielding approximately 6 and 3 times, respectively, the exposure at the MRHD based on summed AUCs of ivacaftor and its metabolites).

### **Use in Pregnancy**

Category B3

Category B3 drugs have been taken by only a limited number of pregnant women and women of childbearing age, without an increase in the frequency of malformation or other direct or indirect harmful effects on the human fetus having been observed. Studies in animals have shown evidence of an increased occurrence of fetal damage, the significance of which is considered uncertain in humans.

There are no adequate and well-controlled studies of KALYDECO in pregnant women. Developmental toxicity studies in animals revealed no teratogenicity in rats at oral doses up to 200 mg/kg/day (yielding 5 times the summed AUC for ivacaftor and its major metabolites anticipated in patients) or in rabbits at up to 100 mg/kg/day (relative exposure based on summed AUCs,  $\geq 3$ ). Fetal weight was decreased and the incidence of minor fetal skeletal abnormalities was increased in rats treated at 200 mg/kg/day; these effects were observed in conjunction with maternal toxicity. Ivacaftor and/or its metabolites were shown to cross the placenta in rats and rabbits. As animal reproduction studies are not always predictive of human response, and given the limited experience with KALYDECO in pregnancy, KALYDECO should only be used during pregnancy if the expected benefits to the mother justify the potential risks to the fetus.

### **Use in Lactation**

It is unknown whether ivacaftor and/or its metabolites are excreted in human milk. Ivacaftor and/or its metabolites were shown to be excreted into the milk of lactating rats. The safe use of KALYDECO during breast-feeding has not been established. KALYDECO should only be used during breast-feeding if the potential benefit outweighs the potential risk.

### **Paediatric Use**

The safety and efficacy of KALYDECO in children aged less than 2 years and weighing less than 10 kg have not been studied. The safety and efficacy of Kalydeco in children aged 2 to less than 6 years beyond 24 weeks have not been studied.

Efficacy was not demonstrated in patients less than 18 years of age with CF who have an *R117H* mutation in clinical trials. Currently available data are described in the Pharmacology, Clinical Trials, Adverse Effects and Dosage and Administration sections.

Cataracts were seen in juvenile rats treated with ivacaftor from postnatal day 7-35 at oral doses  $\geq 10$  mg/kg/day, yielding exposure to ivacaftor and its major metabolites approximately 3.5-6 times lower than that in patients at the MRHD based on summed AUCs. This finding has not been observed in older animals. The potential relevance of these findings in humans is unknown.

### **Cataracts**

Cases of non-congenital lens opacities without impact on vision have been reported in paediatric patients treated with ivacaftor. Although other risk factors were present in some cases (such as corticosteroid use and exposure to radiation), a possible risk attributable to ivacaftor cannot be excluded. Baseline and follow-up ophthalmological examinations are recommended in paediatric patients initiating ivacaftor treatment.

### **Use in the Elderly**

Clinical studies of ivacaftor did not include a sufficient number of patients age 65 years and older to evaluate the efficacy and safety of ivacaftor in this age range. Thus, the efficacy and safety of ivacaftor in elderly patients have not been evaluated.

### **Renal Impairment**

Caution is recommended while using ivacaftor in patients with severe renal impairment (creatinine clearance less than or equal to 30 mL/min) or end-stage renal disease (see DOSAGE AND ADMINISTRATION and PHARMACOLOGY).

### **Hepatic Impairment**

Use of KALYDECO is not recommended in patients with severe hepatic impairment unless the benefits are expected to outweigh the risks of overexposure. In such cases, the starting dose interval should be one tablet or one sachet of KALYDECO every other day (see DOSAGE AND ADMINISTRATION and PHARMACOLOGY).

### **Patients after Organ Transplantation**

KALYDECO has not been studied in patients with CF who have undergone organ transplantation. Therefore, use in transplanted patients is not recommended. See INTERACTIONS WITH OTHER MEDICINES for interactions with ciclosporin or tacrolimus.

### **Lactose**

KALYDECO contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency, or glucose-galactose malabsorption should not take this medicine.

### **Genotoxicity**

Ivacaftor was negative for genotoxicity in the following assays: Ames test for bacterial gene mutation, *in vitro* chromosomal aberration assay in Chinese hamster ovary cells, and *in vivo* mouse micronucleus test.

## **Carcinogenicity**

Two-year oral studies in mice and rats to assess the carcinogenic potential of ivacaftor demonstrated that ivacaftor was not carcinogenic in either species. Plasma exposures to ivacaftor in mice at the non-carcinogenic dosage (200 mg/kg/day, the highest dosage tested) were approximately 4- to 7-fold higher than the plasma levels measured in humans following ivacaftor therapy, and at least 1.2 to 2.4 times higher with respect to the summed AUC for ivacaftor and its major metabolites. Plasma exposures to ivacaftor in rats at the non-carcinogenic dosage (50 mg/kg/day, the highest dosage tested) were approximately 16- to 29-fold higher than the plasma levels measured in humans following ivacaftor therapy, and 6- to 9-fold higher with respect to the summed AUC for ivacaftor and its major metabolites.

## **Effect on Laboratory Tests**

### *Liver Function Tests*

Moderate transaminase (alanine transaminase [ALT] or aspartate transaminase [AST]) elevations are common in patients with CF. In Studies 1, 2, and 3, the incidence and clinical features of transaminase elevations in clinical trials were similar between patients in the ivacaftor and placebo treatment groups (see ADVERSE EFFECTS). In the subset of patients with a medical history of elevated transaminases, increased ALT or AST has been reported more frequently in patients receiving ivacaftor compared to placebo. Therefore, liver function tests are recommended for all patients prior to initiating ivacaftor, every 3 months during the first year of treatment, and annually thereafter. For patients with a history of transaminase elevations, consider more frequent monitoring of liver function tests (see ADVERSE EFFECTS). Patients who develop unexplained increased transaminase levels during treatment should be closely monitored until the abnormalities resolve. Dosing should be interrupted in patients with ALT or AST of greater than 5 times the upper limit of normal (ULN). Following resolution of transaminase elevations, consideration should be given to the continuation of treatment after assessment of the individual benefits and risks.

## **Interactions with Medicinal Products**

Ivacaftor is a substrate of CYP3A4 and CYP3A5 isoenzymes. Medicinal products that inhibit or induce CYP3A activity may impact the pharmacokinetics of ivacaftor. The dose of KALYDECO must be adjusted when concomitantly used with strong and moderate CYP3A inhibitors. Exposure to ivacaftor may be reduced by the concomitant use of CYP3A inducers, potentially resulting in the loss of efficacy of KALYDECO (see INTERACTIONS WITH OTHER MEDICINES).

Ivacaftor is a weak CYP3A and P-glycoprotein (P-gp) inhibitor and may modify the pharmacokinetics of medicinal products that are substrates of CYP3A and/or P-gp. *In vitro* studies indicated that ivacaftor has the potential to inhibit CYP2C9.

## **Driving and Operating Machinery**

Dizziness has been reported in patients receiving KALYDECO, which could influence the ability to drive or operate machines (see ADVERSE EFFECTS). Patients experiencing dizziness should be advised not to drive or operate machines until symptoms abate.

## **INTERACTIONS WITH OTHER MEDICINES**

Ivacaftor is a substrate of CYP3A4 and CYP3A5. It is a weak inhibitor of CYP3A and P-gp and a potential inhibitor of CYP2C9.

## **Effects of Other Medicines on Ivacaftor**

### *CYP3A Inhibitors*

Ivacaftor is a sensitive CYP3A substrate. Co-administration with ketoconazole, a strong CYP3A inhibitor, increased ivacaftor exposure (measured as area under the curve [AUC]) by

8.5-fold and increased hydroxymethyl-ivacaftor (M1) to a lesser extent than ivacaftor. A reduction of the KALYDECO dose to one tablet or one sachet twice a week is recommended for co-administration with strong CYP3A inhibitors, such as ketoconazole, itraconazole, posaconazole, voriconazole, telithromycin, and clarithromycin.

Co-administration with fluconazole, a moderate inhibitor of CYP3A, increased ivacaftor exposure by 3-fold and increased M1 to a lesser extent than ivacaftor. A reduction of the KALYDECO dose to one tablet or one sachet once daily is recommended for patients taking concomitant moderate CYP3A inhibitors, such as fluconazole and erythromycin (see DOSAGE AND ADMINISTRATION).

Co-administration of KALYDECO with grapefruit juice, which contains one or more components that moderately inhibit CYP3A, may increase exposure to ivacaftor. Food containing grapefruit or Seville oranges should be avoided during treatment with KALYDECO.

#### ***CYP3A Inducers***

Co-administration of ivacaftor with rifampicin, a strong CYP3A inducer, decreased ivacaftor exposure (AUC) by 89% and decreased M1 to a lesser extent than ivacaftor.

Co-administration with strong CYP3A inducers, such as rifampicin, rifabutin, phenobarbital, carbamazepine, phenytoin and St. John's wort (*Hypericum perforatum*), is not recommended. Concomitant use of weak to moderate inducers of CYP3A (e.g., dexamethasone, high-dose prednisone) may decrease the exposure of ivacaftor and thus may reduce KALYDECO efficacy.

#### ***Other Recommendations***

Co-administration of ciprofloxacin with ivacaftor did not affect the exposure of ivacaftor. No dose adjustment is required when KALYDECO is co-administered with ciprofloxacin.

### **Effects of Ivacaftor on Other Medicines**

#### ***CYP3A, P-gp, or CYP2C9 Substrates***

Based on *in vitro* results, ivacaftor and its M1 metabolite have the potential to inhibit CYP3A and P-gp. Co-administration with (oral) midazolam, a sensitive CYP3A substrate, increased midazolam exposure 1.5-fold, consistent with weak inhibition of CYP3A by ivacaftor. Co-administration with digoxin, a sensitive P-gp substrate, increased digoxin exposure by 1.3-fold, consistent with weak inhibition by P-gp by ivacaftor. Administration of KALYDECO may increase systemic exposure of medicinal products that are substrates of CYP3A and/or P-gp, which may increase or prolong their therapeutic effect and adverse reactions. Use with caution and monitor for benzodiazepine-related side effects when using concomitant oral midazolam, alprazolam, diazepam, or triazolam. Use with caution and appropriate monitoring when using concomitant digoxin, cyclosporine, or tacrolimus. Ivacaftor may inhibit CYP2C9. Therefore, monitoring of the INR during co-administration with warfarin is recommended.

#### ***Other Recommendations***

Ivacaftor has been studied with an oestrogen/progesterone oral contraceptive and was found to have no significant effect on the exposures of the oral contraceptive. Ivacaftor is not expected to modify the efficacy of oral contraceptives. Therefore, no dose adjustment of oral contraceptives is necessary.

Ivacaftor has been studied with the CYP2C8 substrate rosiglitazone. No significant effect on rosiglitazone exposure was found. Therefore, no dose adjustment of CYP2C8 substrates such as rosiglitazone is necessary.

Ivacaftor has been studied with the CYP2D6 substrate desipramine. No significant effect on desipramine exposure was found. Therefore, no dose adjustment of CYP2D6 substrates such as desipramine is necessary.

Interaction studies have only been performed in adults.

## ADVERSE EFFECTS

### Experience from Clinical Trials

The safety profile of KALYDECO is based on six clinical trials: two pooled, placebo-controlled Phase 3 clinical trials (Studies 1 and 2) conducted in 213 CF patients (109 received ivacaftor and 104 received placebo up to 48 weeks) who had a *G551D* mutation in the *CFTR* gene; a 96-week open-label extension study (Study 4) that included 192 patients with a *G551D* mutation; an 8-week, Phase 3, placebo-controlled crossover design study (Study 5) in 39 patients with CF who had a non-*G551D* gating (class III) mutation in the *CFTR* gene; and a 24-week, placebo-controlled trial (Study 6) involving 69 patients with an *R117H* mutation in the *CFTR* gene; and a 24-week, open-label study (Study 7) that included 34 patients with a *G551D* or another gating (class III) mutation in the *CFTR* gene. Patients treated with KALYDECO in these trials were between the ages of 2 and 68 years.

The most common adverse reactions that were more frequent in patients with a *G551D* mutation who received ivacaftor for 48 weeks in the placebo-controlled Phase 3 trials were abdominal pain (15.6% versus 12.5% on placebo), diarrhoea (12.8% versus 9.6% on placebo), dizziness (9.2% versus 1.0% on placebo), rash (12.8% versus 6.7% on placebo), upper respiratory tract reactions (including upper respiratory tract infection, nasal congestion, pharyngeal erythema, oropharyngeal pain, rhinitis, sinus congestion, and nasopharyngitis) [63.3% versus 50.0% on placebo], headache (23.9% versus 16.3% on placebo) and bacteria in sputum (7.3% versus 3.8% on placebo). One patient in the ivacaftor group reported a serious adverse reaction: abdominal pain.

Adverse events were identified based on the pooled data of the 48-week, placebo-controlled Studies 1 and 2 in patients aged 6 years and older who had a *G551D* mutation in the *CFTR* gene.

Table 7 shows adverse events with an incidence of at least 10% in any treatment group from the two double-blind, placebo-controlled Phase 3 trials.

<b>Preferred Term</b>	<b>KALYDECO N=109 n (%)</b>	<b>Placebo N=104 n (%)</b>
Cystic fibrosis lung	42 (38.5)	58 (55.8)
Cough	40 (36.7)	52 (50.0)
Headache	26 (23.9)	17 (16.3)
Oropharyngeal pain	24 (22.0)	19 (18.3)
Upper respiratory tract infection	24 (22.0)	14 (13.5)
Nasal congestion	22 (20.2)	16 (15.4)
Abdominal pain	17 (15.6)	13 (12.5)
Pyrexia	16 (14.7)	16 (15.4)
Nasopharyngitis	16 (14.7)	12 (11.5)

<b>Preferred Term</b>	<b>KALYDECO N=109 n (%)</b>	<b>Placebo N=104 n (%)</b>
Productive cough	14 (12.8)	16 (15.4)
Diarrhoea	14 (12.8)	10 (9.6)
Rash	14 (12.8)	7 (6.7)
Nausea	13 (11.9)	11 (10.6)
Vomiting	11 (10.1)	17 (16.3)
Rales	11 (10.1)	12 (11.5)
Haemoptysis	9 (8.3)	17 (16.3)
Pulmonary function test decreased	5 (4.6)	15 (14.4)
Abdominal pain upper	10 (9.2)	11 (10.6)

Table 8 presents the adverse reactions identified in patients aged 6 years and older who had a *G551D* mutation in at least one allele listed by system organ class, preferred term, and frequency. Adverse reactions are ranked under the MedDRA frequency classification: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); and not known (frequency cannot be estimated using the available data).

<b>System Organ Class</b>	<b>Frequency Category</b>	<b>Adverse Reactions (Preferred term) KALYDECO N=109</b>
Infections and infestations	very common	Nasopharyngitis
	very common	Upper respiratory tract infection
	common	Rhinitis
Nervous system disorders	very common	Headache
	common	Dizziness
Ear and labyrinth disorders	common	Ear discomfort
	common	Ear pain
	common	Tinnitus
	common	Tympanic membrane hyperaemia
	uncommon	Ear congestion
	uncommon	Vestibular disorder
Respiratory, thoracic, and mediastinal disorders	very common	Nasal congestion
	very common	Oropharyngeal pain
	common	Pharyngeal erythema
	common	Sinus congestion
Gastrointestinal disorders	very common	Abdominal pain
	very common	Diarrhoea
Skin and subcutaneous tissue disorders	very common	Rash

<b>Table 8. Adverse Reactions in KALYDECO-treated Patients Age 6 Years and Older with the G551D Mutation in the CFTR Gene</b>		
<b>System Organ Class</b>	<b>Frequency Category</b>	<b>Adverse Reactions (Preferred term) KALYDECO N=109</b>
Reproductive system and breast disorders	uncommon	Breast inflammation
	uncommon	Breast mass
	uncommon	Gynaecomastia
	uncommon	Nipple disorder
	uncommon	Nipple pain
Investigations	common	Bacteria in sputum

## **Description of Selected Adverse Reactions**

### ***Rash***

During 48-week, placebo-controlled clinical trials, the incidence of rash was 12.8% in KALYDECO-treated patients. Including data from all clinical trial and post-marketing data, most of these events were non-serious and most of these patients did not discontinue the treatment because of rash.

### ***Ear and Labyrinth Disorders***

During 48-week, placebo-controlled clinical trials, the incidence of ear and labyrinth disorders was 9.2% in KALYDECO-treated patients. Most events were described as mild to moderate in severity, 1 event of ear pain was described as severe, none were serious, and no patients discontinued treatment because of ear and labyrinth disorders.

### ***Nervous System Disorders***

#### ***Headache***

During 48-week, placebo-controlled clinical trials, the incidence of headache was 23.9% in KALYDECO-treated patients. Including data from all clinical trial and post-marketing data, most of these events were non-serious and most of these patients did not discontinue the treatment because of headache.

#### ***Dizziness***

During 48-week, placebo-controlled clinical trials, the incidence of dizziness was 9.2% in the KALYDECO-treated patients. Including data from all clinical trial and post-marketing data, most of these events were non-serious and most of these patients did not discontinue the treatment because of dizziness.

### ***Upper Respiratory Tract Reactions***

During 48-week, placebo-controlled clinical trials, the incidence of upper respiratory tract reactions (upper respiratory tract infection, nasal congestion, pharyngeal erythema, oropharyngeal pain, rhinitis, sinus congestion, and nasopharyngitis) was 63.3% in KALYDECO-treated patients. Most events were described as mild to moderate in severity, 1 event of upper respiratory tract infection and 1 event of nasal congestion were considered to be severe, none were serious, and no patients discontinued treatment because of upper respiratory tract reactions.

### ***Laboratory Abnormalities***

#### ***Transaminase Elevations***

During the placebo-controlled Studies 1, 2, and 3, up to 48 weeks, in patients with a G551D mutation in the CFTR gene or who were homozygous for the F508del mutation, the incidence of maximum transaminase (ALT or AST) >8, >5 or >3 x ULN was 1.8%, 2.3%, and 5.9% in

KALYDECO-treated patients and 1.5%, 2.3%, and 8.3% in placebo-treated patients, respectively. Three patients, 2 (1.5%) on placebo and 1 (0.5%) on KALYDECO, permanently discontinued treatment for elevated transaminases, which were all >8 x ULN. No KALYDECO-treated patients experienced a transaminase elevation >3 x ULN associated with elevated total bilirubin >1.5 x ULN. In KALYDECO-treated patients, most transaminase elevations up to 5 x ULN resolved without treatment interruption. KALYDECO dosing was interrupted in most patients with transaminase elevations >5 x ULN. In all instances where dosing was interrupted for elevated transaminases, KALYDECO dosing was able to be resumed (see PRECAUTIONS).

### **Paediatric Population**

The safety data were evaluated in 34 patients between 2 to less than 6 years of age, 52 patients between 6 to less than 12 years of age and 93 patients between 12 to less than 18 years of age.

The safety profile is generally consistent among children and adolescents and is also consistent with adult patients.

### **Long-term Safety (Study 4)**

Among 192 patients with the *G551D* mutation treated with KALYDECO for 96 weeks in an open-label extension study (Study 4) following the placebo-controlled Phase 3 studies (96 to 144 weeks cumulatively), the nature of adverse events was similar to those reported in the placebo-controlled Phase 3 studies. Serious adverse reactions observed during the 96-week extension study included abdominal pain (1%), headache (1%), and vestibular disorder (0.5%).

### **Non-*G551D* Gating Population (Study 5)**

In an 8-week, two-part, randomised, double-blind, placebo-controlled, crossover Phase 3 clinical trial of 39 patients with CF aged 6 and older who had a non-*G551D* gating mutation in the *CFTR* gene (Study 5), the safety results were consistent with those observed in studies in patients with CF who had the *G551D* mutation. In the non-*G551D* gating population, one adverse reaction occurred in more patients during treatment with KALYDECO, compared with placebo: rhinitis (7.9% versus 5.4% on placebo).

### **R117H Population (Study 6)**

In a 24-week, placebo-controlled trial (Study 6) involving 69 patients with CF aged 6 and older who had an *R117H* mutation in the *CFTR* gene, the safety results were consistent with those observed in studies in patients with CF who had the *G551D* mutation.

### **Paediatric Safety and Efficacy Study (Study 7)**

In a 24-week, open-label study in 34 patients 2 to less than 6 years of age with CF who had a *G551D* or another gating (class III) mutation in the *CFTR* gene (Study 7), the safety results were consistent with those observed in patients with CF who had the *G551D* mutation. In the paediatric population transaminase elevation and vomiting were observed more frequently as compared to the adults; however, it is unknown if these effects were due to Kalydeco. In this study, the incidence of patients experiencing transaminase elevations (ALT or AST) >3 x ULN was 14.7% (5/34). All 5 patients had maximum ALT or AST levels >8 x ULN, which returned to baseline levels following interruption of ivacaftor dosing. Ivacaftor was discontinued permanently in one patient (see PRECAUTIONS for management of elevated transaminases).

### **Post-Marketing Experience**

There are no relevant updates from the post-marketing experience.

## DOSAGE AND ADMINISTRATION

**KALYDECO should only be prescribed by physicians with experience in the treatment of cystic fibrosis.** If the patient's genotype is unknown, an accurate and validated genotyping method should be performed to confirm the presence of a gating (class III) mutation or an *R117H* mutation in at least one allele of the *CFTR* gene before starting treatment.

In patients with *R117H* mutation, analysis of the poly-T variants may be considered to better define those more likely to respond to treatment.

*Adults, adolescents, and children aged 2 years and older should be dosed according to Table 9.*

<b>Weight</b>	<b>Dose</b>	<b>Total daily dose</b>
<14 kg	50 mg granules (one sachet) q12h	100 mg (two sachets)
≥14 kg to <25 kg	75 mg granules (one sachet) q12h	150 mg (two sachets)
≥25 kg	150 mg tablet q12h	300 mg (two tablets)

KALYDECO should be taken with a fat-containing meal or snack. Meals and snacks recommended in CF guidelines or meals recommended in standard nutritional guidelines contain adequate amounts of fat. A serving size of foods appropriate for age from a typical CF diet should be given. Examples of meals that contain fat are those prepared with butter or oils or those containing eggs, cheeses, nuts, avocados, whole milk, full-fat yoghurt, or meats. Food containing grapefruit or Seville oranges should be avoided during treatment with KALYDECO (see INTERACTIONS WITH OTHER MEDICINES).

### **Use in the Elderly**

The efficacy and safety of KALYDECO in patients age 65 years or older have not been evaluated.

### **Use in Renal Insufficiency**

No dose adjustment is necessary for patients with mild to moderate renal impairment. Caution is recommended while using ivacaftor in patients with severe renal impairment (creatinine clearance less than or equal to 30 mL/min) or end-stage renal disease (see PRECAUTIONS and PHARMACOLOGY).

### **Use in Hepatic Insufficiency**

No dose adjustment is necessary for patients with mild hepatic impairment (Child-Pugh Class A). A reduced dose of KALYDECO in patients with moderate hepatic impairment (Child-Pugh Class B) is recommended as follows: in patients with body weight less than 14 kg, one 50 mg sachet of granules once daily; in patients with body weight 14 kg to less than 25 kg, one 75 mg sachet of granules once daily; and in patients with body weight 25 kg or greater, one 150 mg tablet once daily. There is no experience of the use of KALYDECO in patients with severe hepatic impairment. The use of KALYDECO in these patients is therefore not recommended unless the benefits outweigh the risks. In such cases, the starting dose of KALYDECO should be as follows: in patients with body weight less than 14 kg, one 50 mg sachet of granules every other day; in patients with body weight 14 kg to less than 25 kg, one 75 mg sachet of granules every other day; and in patients with body weight 25 kg or greater, one 150 mg tablet every other day. Dosing intervals should be modified according to clinical response and tolerability (see DOSAGE AND ADMINISTRATION, PRECAUTIONS and PHARMACOLOGY).

### **Use in Combination with Other Medicinal Compounds**

When co-administered with strong inhibitors of CYP3A (e.g., ketoconazole, itraconazole, posaconazole, voriconazole, telithromycin, and clarithromycin), KALYDECO should be administered as follows: in patients with body weight less than 14 kg, one 50 mg sachet of granules twice a week; in patients with body weight 14 kg to less than 25 kg, one 75 mg sachet of granules twice a week; and in patients with body weight 25 kg or greater, one 150 mg tablet twice a week.

When co-administered with moderate inhibitors of CYP3A (e.g., fluconazole, erythromycin), KALYDECO should be administered as follows: in patients with body weight less than 14 kg, one 50 mg sachet of granules once daily; in patients with body weight 14 kg to less than 25 kg, one 75 mg sachet of granules once daily; and in patients with body weight 25 kg or greater, one 150 mg tablet once daily (see DOSAGE AND ADMINISTRATION, PRECAUTIONS and INTERACTIONS WITH OTHER MEDICINES).

### **Method of Administration**

For oral use. Patients should be instructed to swallow the tablets whole (i.e., patients should not chew, break, or dissolve the tablet).

#### *Tablets*

Patients should be instructed to swallow the tablets whole (i.e., patients should not chew, break or dissolve the tablet).

#### *Granules*

The entire content of each sachet of granules should be mixed with one teaspoon (5 mL) of age-appropriate soft food or liquid and the mixture completely consumed. Food or liquid should be at room temperature or below. Each sachet is for single use only. Once mixed, the product has been shown to be stable for one hour, and therefore should be ingested during this period. Some examples of soft foods or liquids include puréed fruits or vegetables, yogurt, applesauce, water, milk, or juice. A fat-containing meal or snack should be consumed just before or just after dosing.

### **OVERDOSAGE**

For information on the management of overdose, contact the Poisons Information Centre on 131126 (Australia).

The highest repeated dose was 450 mg (in a tablet formulation) every 12 hours for 4.5 days (9 doses) in a trial evaluating the effect of KALYDECO on ECGs in healthy subjects.

Adverse events reported at a higher incidence compared to placebo included dizziness and diarrhoea.

No specific antidote is available for overdose with KALYDECO. Treatment of overdose with KALYDECO consists of general supportive measures including monitoring of vital signs, liver function tests, and observation of the clinical status of the patient.

### **PRESENTATION AND STORAGE CONDITIONS**

#### *Tablets*

KALYDECO (ivacaftor) film-coated tablets are supplied as light blue, capsule-shaped tablets (16.5 mm x 8.4 mm in modified caplet shape) containing 150 mg of ivacaftor. Each tablet is printed with “V 150” in black ink on one side only.

KALYDECO tablets are packaged in a blister pack or a child-resistant bottle with desiccant.

The tablets are packaged in a thermoform (polychlorotrifluoroethylene [PCTFE]/foil) blister or a high-density polyethylene (HDPE) bottle with a polypropylene, foil-lined induction seal closure and molecular sieve desiccant.

The following pack sizes are available:

- Blister pack containing 56 film-coated tablets
- Bottle containing 56 film-coated tablets

#### *Granules*

KALYDECO (ivacaftor) granules are supplied as white to off-white sweetened, unflavoured granules approximately 2 mm in diameter enclosed in unit dose packets containing 50 mg or 75 mg of ivacaftor per sachet.

Granules are packaged in a Biaxially Oriented Polyethylene Terephthalate/Polyethylene/Foil/Polyethylene (BOPET/PE/Foil/PE) sachet.

Pack size of 56 sachets (contains 4 individual wallets with 14 sachets per wallet) is available.

Store below 30°C.

#### **NAME AND ADDRESS OF THE SPONSOR**

Vertex Pharmaceuticals (Australia) Pty Ltd  
Suite 3, Level 3  
601 Pacific Highway  
St Leonards  
NSW 2065  
Australia

#### **POISON SCHEDULE OF THE MEDICINE**

S4 - Prescription only medicine

#### **DATE OF FIRST INCLUSION IN THE AUSTRALIAN REGISTER OF THERAPEUTIC GOODS**

**09 July 2013**

#### **DATE OF MOST RECENT AMENDMENT**

**16 September 2016**

© 2016 Vertex Pharmaceuticals Incorporated

KALYDECO, VERTEX and the VERTEX triangle logo are registered trademarks of Vertex Pharmaceuticals Incorporated.