

# **AUSTRALIAN PRODUCT INFORMATION – APOMINE<sup>®</sup> INTERMITTENT (Apomorphine hydrochloride hemihydrate) INJECTION CARTRIDGE**

## **1. NAME OF THE MEDICINE**

Apomorphine hydrochloride hemihydrate

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Apomine Intermittent is a sterile solution for injection containing 10 mg/mL of apomorphine hydrochloride hemihydrate in Water for Injections BP. Each 3 mL cartridge contains 30 mg apomorphine hydrochloride hemihydrate. Sodium metabisulfite 1 mg/mL is included in the formulation as an antioxidant.

Excipient(s) with known effect:

Sodium metabisulfite

For the full list of excipients, see Section 6.1 List of excipients.

## **3. PHARMACEUTICAL FORM**

Apomine Intermittent is a clear, colourless to slightly yellow sterile solution for subcutaneous injection, free from visible particles.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Apomine Intermittent is indicated to reduce the number and severity of 'off' phases in patients with Parkinson's disease severely disabled by motor fluctuations refractory to conventional therapy. Initiation of therapy with Apomine Intermittent should be undertaken in a specialist unit in a hospital setting. Conventional therapy should be continued during 'on' phases.

### **4.2 Dose and method of administration**

#### **Dosage**

The optimal dosage of Apomine Intermittent has to be determined on an individual patient basis. Hospital admission under appropriate specialist supervision is advised when establishing a patient's therapeutic regimen.

It is essential that the patient is established on the antiemetic domperidone for at least 48 - 72 hours prior to initiation of therapy.

**Patient selection:** For patients in whom conventional therapy has failed, Apomine Intermittent injections are only considered to be suitable for Parkinson's disease patients capable of recognising and anticipating 'off' phases in motor performance. Patients must be capable and motivated for Apomine Intermittent to be used effectively. Adult patients through all age ranges have been successfully managed with apomorphine injections. Apomine Intermittent is contraindicated in children and adolescents below 18 years of age.

Elderly patients, in appropriate circumstances, can be successfully managed with Apomine Intermittent.

The practical steps described below should be followed when commencing a patient on treatment:

- Pretreat with domperidone.
- Discontinue all existing antiparkinsonian medication to provoke an 'off' phase in motor performance.
- Determine the threshold dose response to Apomine Intermittent that produces an unequivocal motor response.
- Re-establish other antiparkinsonian agents.
- Determine effective treatment regimen for Apomine Intermittent.
- Teach patient and/or carer how and when to administer.
- Discharge from hospital.
- Monitor frequently and adjust dosage regimen as appropriate.
- Full details are given below.

**Pretreatment:** Domperidone is a peripherally acting dopamine receptor antagonist given by mouth to prevent nausea and vomiting. Domperidone is commenced 48 - 72 hours prior to the first dose of Apomine Intermittent. When patients are stabilised with respect to dosage of Apomine Intermittent, the dose of domperidone is reduced by 10 mg per day every week until mild nausea appears. The maintenance dose of domperidone is the lowest level which completely prevents nausea. Domperidone can usually be withdrawn after several weeks. Before the decision to initiate domperidone and apomorphine treatment, risk factors for QT interval prolongation in the individual patient should be carefully assessed to ensure that the benefit outweighs the risk (see Section 4.4 Special warnings and precautions for use). The cardiovascular assessment should include an ECG and QT measurement. Patients with severe renal insufficiency will require the dosing interval of domperidone to be changed from three times a day to once or twice a day. For further information regarding domperidone refer to the domperidone product information..

**Provoking and assessing an 'off' phase:** After at least 3 days of hospitalisation, all antiparkinsonian therapy is withheld overnight to provoke an 'off' phase in motor performance and to undertake a baseline motor assessment as follows:

- a) Alternate, unilateral hand tapping for 30 seconds on mounted digital counters (preferably 20 cm apart).
- b) Time taken to walk 12 metres.
- c) Clinical assessment of tremor and dyskinesia according to a four point scale (0 = nil, 1 = mild, 2 = moderate, 3 = severe).
- d) Scoring on a modified Webster disability scale to assess 12 features of parkinsonism (maximum disability score of 36).

**Determination of the threshold dose:** Following baseline motor assessment, the patient is challenged for Apomine Intermittent responsiveness according to the following schedule:

- 1.5 mg Apomine Intermittent (0.15 mL) is injected subcutaneously (SC) and the patient is observed over 30 minutes for motor responsiveness.
- If no or poor response is obtained, a second dose of 3 mg Apomine Intermittent (0.3 mL) is given 40 minutes after the first dose, and the patient observed for a further 30 minutes.
- The dosage is increased in an incremental fashion every 40 minutes and the patient observed carefully for an unequivocal motor response. The third dose is 5 mg SC, and the fourth dose is 7 mg SC. If the patient shows no response to the 7 mg dose then the patient must be classified as a non-responder to Apomine Intermittent and no further attempts to provoke a motor response should be made. If the patient shows only a mild response to the 7 mg dose, a maximum dose of 10 mg can be used to see if an unequivocal motor response is possible.
- The lowest dose producing an unequivocal motor response is called the threshold dose. For the majority of patients, the threshold dose is less than 7 mg of Apomine Intermittent (0.7 mL), although very occasionally it can be up to 10 mg of Apomine Intermittent (1.0 mL).

Motor responsiveness is judged to be positive if 2 or more of the following are seen:

- a) More than 15% increase in tapping score.
- b) More than 25% improvement in walking time.
- c) An improvement of at least 2 points of tremor score.
- d) An improvement of Webster's score of 3 or more.

**Initiation of treatment:** Following establishment of an acceptable threshold dose of Apomine Intermittent, the patient should be restarted on conventional antiparkinsonian therapy.

A subcutaneous injection of the established threshold dose may then be given into the lower abdomen or outer thigh at the first signs of an 'off' phase. The patient should then be observed over the following hour and the quality of their 'on' phase noted. It may be appropriate to modify the dose of Apomine Intermittent according to the patient's response.

Close monitoring of therapeutic benefits and adverse reactions under specialist supervision is required after initiation of treatment.

### **Method of Administration**

Apomine Intermittent is administered by subcutaneous injection. Apomine Intermittent injection is either into the anterior abdominal wall or anterolateral thigh. The usual dosage range is 2.4 to 3.6 mg per injection; the maximum single dose being 6 mg and the maximum total daily dose being 50 mg.

To ensure accurate dosing, either 1.0 mL insulin syringes or a D-mine<sup>®</sup> Pen should be used to administer Apomine Intermittent injections. The injection is given in an undiluted form.

Apomine Intermittent must not be used via the intravenous route.

Chemical and physical in-use stability has been demonstrated for 15 days at 25°C. However, as the product contains no antimicrobial agent, for microbial reasons, once opened the contents of the cartridge should be used within 72 hours.

Do not use if the solution has turned green. The solution should be inspected visually prior to use. Only clear, colourless to slightly yellow and particle-free solution should be used (also see section 6.4 Special precautions for storage).

Patients who have shown a good 'on' phase response during the initiation stage, but whose overall control remains unsatisfactory using intermittent injections, or who require many and frequent injections (e.g. 8-10 injections per day), may be commenced on or transferred to continuous subcutaneous infusion by minipump. For patients requiring continuous subcutaneous infusion, Apomine Solution for Infusion should be considered.

**Monitoring treatment:** Long term specialist supervision of patients is advised.

There is a high probability of adverse effects to Apomine Intermittent therapy. The frequency and severity of adverse events should be monitored carefully at regular intervals and a reassessment of the patient carried out if appropriate. Adjustments to the dosage or discontinuation may be necessary.

### **4.3 Contraindications**

Apomine Intermittent is contraindicated in patients with a known hypersensitivity or allergy to apomorphine, morphine or to any of the excipients (including sodium metabisulfite) or chemically related products.

Apomine Intermittent should not be administered to patients with pre-existing neuropsychiatric problems or dementias due to either pathological processes, e.g. Alzheimer's disease, or to patients whose 'on' response to L-dopa is marred by severe dyskinesia, hypotonia or psychotoxicity.

Apomorphine is also contraindicated in patients with inadequate renal or liver function, unstable coronary vascular disease, cerebrovascular disease, respiratory depression or CNS depression.

Apomine Intermittent is contraindicated for children and adolescents under 18 years of age. Apomine Intermittent is also contraindicated in patients with a known hypersensitivity to sodium metabisulfite.

### **4.4 Special warnings and precautions for use**

For Subcutaneous Use Only (see Section 4.8 Adverse effects (Undesirable effects)).

Patients sensitive to morphine or its derivatives may be sensitive to apomorphine. Apomorphine should therefore not be administered to patients with a known hypersensitivity or allergy to apomorphine, morphine or chemically related compounds (see Section 4.3 Contraindications).

Apomine Intermittent contains sodium metabisulfite which may cause allergic type reactions, including anaphylactic symptoms and life threatening or less severe asthmatic episodes in susceptible people.

In patients with cardiac decompensation or cerebrovascular disease, vomiting may cause an increase in blood pressure that may lead to haemorrhage and vascular accidents. Apomorphine is therefore contraindicated in these patients (see Section 4.3 Contraindications).

Caution should be used in administering apomorphine to patients with a predisposition to nausea and vomiting. Apomorphine may cause an increased risk of persistent vomiting. A risk-benefit assessment should be considered in these patients.

Since apomorphine may produce hypotension, even when given with domperidone pre-treatment, care should be exercised in patients with pre-existing cardiac disease or in patients taking vasoactive medicinal products such as anti-hypertensives, and especially in patients with pre-existing postural hypotension.

Since apomorphine, especially at high doses, may have the potential for QT prolongation, caution should be exercised when treating patients at risk for Torsades de pointes arrhythmia.

When used in combination with domperidone, risk factors in the individual patient should be carefully assessed. This should be done before treatment initiation, and during treatment. Important risk factors include serious underlying heart conditions such as congestive cardiac failure, severe hepatic impairment or significant electrolyte disturbance. Also, medication possibly affecting electrolyte balance, CYP3A4 metabolism or QT interval should be assessed.

Monitoring for an effect on the QTc interval is advisable. An ECG should be performed:

- prior to treatment with domperidone
- during the treatment initiation phase
- as clinically indicated thereafter.

The patient should be instructed to report possible cardiac symptoms including palpitations, syncope, or near-syncope. They should also report clinical changes that could lead to hypokalaemia, such as gastroenteritis or the initiation of diuretic therapy. At each medical visit, risk factors should be revisited.

Apomorphine is associated with local subcutaneous effects. These can sometimes be reduced by the rotation of injection sites or possibly by the use of ultrasound (if available) in order to avoid areas of nodularity and induration (see Section 4.8 Adverse effects (undesirable effects)).

Caution is advised when combining apomorphine with other medicinal products, especially those with a narrow therapeutic range (see Section 4.5 Interactions with other medicines and other forms of interactions).

Neuropsychiatric problems co-exist in many patients with advanced Parkinson's disease. There is evidence that for some patients, neuropsychiatric disturbances may be exacerbated by apomorphine. Special care should be exercised when apomorphine is used in these patients.

The use of apomorphine in conjunction with levodopa treatment may cause Coombs' positive haemolytic anaemia. An initial screen prior to commencement of treatment and at 6 monthly intervals is recommended. In the event of the development of a haemolytic anaemia, a haematological specialist should be consulted. The dose of apomorphine and/or levodopa should be reduced, with careful monitoring of the patient's motor state. It may be necessary to discontinue treatment with levodopa and/or apomorphine in the event that it is not possible to control the anaemia satisfactorily.

Apomorphine has been associated with somnolence, and other dopamine agonists can be associated with sudden sleep onset episodes, particularly in patients with Parkinson's disease. Patients must be informed of this and advised to exercise caution while driving or operating machines during treatment with apomorphine. Patients who have experienced somnolence, and/or an episode of sudden sleep onset, must refrain from driving or operating machines. Furthermore, a reduction of dosage or termination of therapy may be considered.

Apomorphine should be used with caution in patients with endocrine, renal, pulmonary or cardiovascular disease.

Periodic evaluation of hepatic, haemopoietic, renal and cardiovascular function is advised.

Patients with severe renal insufficiency may require the dosing interval for domperidone to be less frequent (see Section 4.2 Dose and method of administration – Pretreatment).

#### Compulsive behaviour

Patients should be regularly monitored for the development of impulse control disorders. Patients and carers should be made aware that behavioural symptoms of impulse control disorders including pathological gambling, increased libido, hypersexuality, compulsive spending or buying, binge eating and compulsive eating can occur in patients treated with dopamine agonists including apomorphine. Dose reduction/tapered discontinuation should be considered if such symptoms develop.

Dopamine Dysregulation Syndrome (DDS) is an addictive disorder resulting in excessive use of apomorphine, seen in some patients treated with apomorphine. Before initiation of treatment, patients and caregivers should be warned of the potential risk of developing DDS.

#### **Use in debilitated patients**

Extra caution is also recommended in debilitated patients, since they may show an increased susceptibility or be more sensitive to the respiratory depressant effects of apomorphine.

#### **Use in the elderly**

Caution is recommended in debilitated or geriatric patients, since they may show an increased susceptibility or be more sensitive to the respiratory depressant effects of apomorphine. Extra caution is recommended during initiation of therapy in elderly patients because of the risk of postural hypotension.

#### **Paediatric use**

Apomine Intermittent is contraindicated for children and adolescents under 18 years of age.

#### **Effects on laboratory tests**

Positive Coombs' tests have been reported for patients receiving apomorphine.

### **4.5 Interactions with other medicines and other forms of interactions**

Patients selected for treatment with apomorphine are almost certain to be taking concomitant medications for their Parkinson's disease. In the initial stages of apomorphine therapy the patient should be monitored for unusual side-effects or signs of potentiation of effect.

Drugs which interfere with central amine mechanisms such as tetrabenazine, metoclopramide, antipsychotic dopamine blocking agents (such as phenothiazines, thioxanthenes and butyrophenones), amphetamines and papaverine should be avoided. If their administration is considered essential, extreme care should be taken and the patient monitored for signs of potentiation, antagonism or other interactions and for any unusual adverse effects.

Neuroleptic medicinal products may have an antagonistic effect if used with apomorphine. There is a potential interaction between clozapine and apomorphine.

The possible side effects of apomorphine on the plasma concentrations of other medicinal products have not yet been studied. Therefore, caution is advised when combining apomorphine with other medicinal products, especially those with a narrow therapeutic range.

#### Antihypertensive and cardiac active medicinal drugs

Even when co-administered with domperidone, apomorphine may potentiate the antihypertensive effects of antihypertensive and cardiac active medicinal products.

It is recommended to avoid the administration of apomorphine with other drugs known to prolong the QT interval.

## **4.6 Fertility, pregnancy and lactation**

### **Effects on fertility**

In a fertility study in male rats, fertility was decreased at 2 mg/kg/day SC, one tenth that of the maximum recommended human dose (based on body surface area). Effects on female fertility have not been determined.

### **Use in pregnancy – Pregnancy Category B3<sup>1</sup>**

The safety of using apomorphine during pregnancy has not been established in either human or animal studies. Apomorphine should therefore not be used in pregnant women, or those likely to become pregnant.

---

<sup>1</sup> Category B3: Drugs which have been taken by only a limited number of pregnant women and women of childbearing age, without an increase in the frequency of malformation or other direct or indirect harmful effects in the human foetus having been observed. Studies in animals have shown evidence of an increased occurrence of foetal damage the significance of which is considered uncertain in humans.

### **Use in lactation**

It is not known whether apomorphine is excreted in breast milk although problems in humans have not been documented. Nevertheless, because many drugs are excreted in human milk and because of the potential for serious adverse drug reactions due to apomorphine in breastfed infants, a decision should be made either to discontinue breastfeeding or the drug, taking into account the importance of the drug to the mother.

## **4.7 Effects on ability to drive and use machines**

Apomorphine has varying degrees of impairment which influences the ability to drive and use machines.

Patients being treated with apomorphine and presenting with somnolence must be informed to refrain from driving or engaging in activities where impaired alertness may put themselves or others at risk of serious injury or death (e.g. operating machines) unless patients have overcome such experiences of somnolence (see also Section 4.4 Special warnings and precautions for use).

## **4.8 Adverse effects (undesirable effects)**

### *Very Common (>10%)*

Most patients experience injection site reactions, particularly with continuous use. These may include subcutaneous nodules, induration (see Section 4.4 Special warnings and precautions for use), erythema, tenderness and panniculitis. Various other local reactions (such as irritation, itching, bruising, fibrosis and pain) may also occur (see Section 4.4 Special warnings and precautions for use). Care should be taken to ensure that areas of ulceration do not become infected.

Hallucinations have been reported.

### *Common (1-10%)*

Gastrointestinal side effects including nausea and vomiting appear to be the most prevalent adverse effects, however tolerance to these effects develops rapidly. Pretreatment with domperidone may reduce or prevent these effects.

Transient sedation with each dose of apomorphine at the start of therapy may occur; this usually resolves over the first few weeks.

Apomorphine is associated with somnolence. Drowsiness and sedation occur in most patients on initial treatment but these effects largely subside with repeated dosing, although in some patients these effects may persist. Tachyphylaxis to postural related faintness or syncope also occurs rapidly.

Neuropsychiatric disturbances are common in parkinsonian patients. Apomorphine should be used with special caution in these patients. Neuropsychiatric disturbances (including transient mild confusion and visual hallucinations) have occurred during apomorphine therapy.

Yawning has been reported during apomorphine therapy.

### *Uncommon (0.1- 1%)*

Apomorphine may induce dyskinesias during 'on' periods, which can be severe in some cases, and in a few patients may result in cessation of therapy. Apomorphine has been associated with sudden sleep onset episodes (see Section 4.4 Special warnings and precautions for use). Postural hypotension is seen infrequently and is usually transient (See section 4.4 Special warnings and precautions for use).

Breathing difficulties have been reported.

Local and generalised rashes have been reported. Injection site skin necrosis and ulceration has been reported.

Haemolytic anaemia and thrombocytopenia have been reported in patients treated with apomorphine (see Section 4.4 Special warnings and precautions for use).

*Rare (0.01 – 0.1%)*

Eosinophilia has rarely occurred during treatment with apomorphine. Peripheral blood eosinophilia, elevated by up to 10%, has occurred in patients on continuous subcutaneous infusion of apomorphine. Blood counts returned to normal in about half of the patients who received treatment over one year.

Due to the presence of sodium metabisulfite, allergic reactions (including anaphylaxis and bronchospasm) may occur.

#### *Other*

Pathological gambling, increased libido, hypersexuality, compulsive spending or buying, binge eating and compulsive eating can occur in patients treated with dopamine agonists including apomorphine (see Section 4.4 Special warnings and precautions for use).

Aggression and agitation have also been reported.

Headache has been reported.

Peripheral oedema has been reported.

Other adverse reactions to apomorphine that have been reported infrequently include stomatitis, confusion, transient rises in serum prolactin, transient metallic taste, spontaneous penile erection, rhinorrhoea, increased lacrimation, reduced facial hair growth and loss of libido.

### **Reporting suspected adverse effects**

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems).

## **4.9 Overdose**

### **Symptoms**

The clinical features of overdose of Apomine Intermittent are an extension of the pharmacological effects of the drug. They include nausea and persistent vomiting, dyskinesias, hypotension and acute circulatory failure, cardiac arrest, respiratory depression, drowsiness and central nervous system depression or stimulation, euphoria, restlessness and hallucinations and possibly coma and death. Concomitant use of domperidone may exacerbate the clinical features of overdose.

### **Treatment**

An opioid antagonist such as naloxone may be given to treat excessive vomiting, central nervous system depression and respiratory depression due to Apomine Intermittent overdose. Excessive vomiting may also be treated with domperidone. Atropine may be also used to treat bradycardia. To treat hypotension, appropriate measures should be taken.

For information on the management of overdose, contact the Poisons Information Centre on 13 11 26 (Australia).

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

#### Mechanism of action

Apomorphine is a directly acting dopamine receptor agonist, structurally related to dopamine. Apomorphine has high *in vitro* binding affinity for the dopamine D<sub>4</sub> and D<sub>5</sub> receptor (*K<sub>i</sub>*: 4 and 14 nM respectively), moderate affinity (*K<sub>i</sub>*: 26 to 130 nM) for the dopamine D<sub>2</sub> and D<sub>3</sub>, adrenergic  $\alpha_{1D}$ ,  $\alpha_{2B}$ ,  $\alpha_{2C}$  receptors, serotonin 5HT<sub>1A</sub>, 5HT<sub>2A</sub>, 5HT<sub>2B</sub>, and 5HT<sub>2C</sub> receptors and low affinity for the dopamine D<sub>1</sub> receptor (*K<sub>i</sub>*: 370 nM). Apomorphine exhibits no affinity for the adrenergic  $\beta_1$  and  $\beta_2$  or histamine H<sub>1</sub> receptors.

The effect of apomorphine as an antiparkinsonian agent is believed to be the result of direct stimulation of postsynaptic D<sub>2</sub> dopamine receptors, but stimulation of presynaptic D<sub>2</sub> dopamine receptors and antagonism of  $\alpha_2$  adrenergic receptors may also be important. Apomorphine reduces the tremor, rigidity and bradykinesia in patients receiving levodopa. Apomorphine induces vomiting by direct stimulation of the medullary chemoreceptor trigger zone.

#### Clinical trials

No data available.

### 5.2 Pharmacokinetic properties

The peripheral pharmacokinetics of apomorphine have been studied following subcutaneous injection, subcutaneous infusion and intravenous infusion.

#### Absorption

Following intramuscular or subcutaneous administration, apomorphine is reported to be well absorbed. Peak plasma concentration occurs as early as three minutes following subcutaneous bolus injection. The rapid and complete absorption from subcutaneous tissues and rapid clearance is believed to correlate with the rapid onset and brief duration of action respectively. Antiparkinsonian effects are observed within five minutes following subcutaneous bolus administration.

#### Distribution

The distribution half-life of apomorphine was found to be five minutes. The volume of distribution, plasma clearance and half-life were similar for subcutaneous injection, subcutaneous infusion and intravenous infusion.

Apomorphine reaches a concentration in the brain up to eight times higher than that in plasma, due to high lipid solubility which allows rapid equilibration between blood and tissue compartments.

#### Metabolism

Apomorphine is metabolised in the liver. Routes of metabolism in humans include sulfation, N-demethylation, glucuronidation and oxidation to norapomorphine by CYP 2B6, CYP 2C8 and CYP 3A4. The major metabolite in humans after sublingual administration was apomorphine sulfate.

## **Excretion**

Apomorphine is cleared rapidly. The elimination half-life ( $t_{1/2}$ ) is about 33 minutes.

## **5.3 Preclinical safety data**

### **Genotoxicity**

*In vitro* genotoxicity studies demonstrated mutagenic and clastogenic effects, most likely due to products formed by oxidation of apomorphine. Apomorphine was not genotoxic *in vivo* in a mouse micronucleus test or in a rat unscheduled DNA synthesis test.

### **Carcinogenicity**

No carcinogenicity studies have been performed.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sodium metabisulfite

Water for injections

Hydrochloric acid

Sodium hydroxide

### **6.2 Incompatibilities**

Incompatibilities were either not assessed or not identified as part of the registration of this medicine. See also Section 4.5 Interactions with other medicines and other forms of interactions.

### **6.3 Shelf life**

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

Chemical and physical in-use stability has been demonstrated for 15 days at 25°C. However, as the product contains no antimicrobial agent, for microbial reasons, once opened the contents of the cartridge should be used within 72 hours.

### **6.4 Special precautions for storage**

Store below 25 °C. Do not refrigerate or freeze.

Protect from light. (Keep the container in the outer carton).

### **6.5 Nature and contents of container**

Clear type I glass cartridges, with bromobutyl rubber stopper and an aluminium cap with bromobutyl rubber seal.

30 mg/3 mL cartridge. Pack of 5 cartridges.

## 6.6 Special precautions for disposal


In Australia, any unused medicine or waste material should be disposed of in accordance with local requirements.

## 6.7 Physicochemical properties

The pH of the injection is 2.5 to 4.0.

### Chemical structure

The chemical structure is shown below:



### CAS number

The CAS registry number of apomorphine hydrochloride hemihydrate is 41372-20-7.

## 7. MEDICINE SCHEDULE (POISONS STANDARD)

Schedule 4

## 8. SPONSOR

Pfizer Australia Pty Ltd  
Level 17, 151 Clarence Street  
Sydney NSW 2000  
Toll Free Number: 1800 675 229  
[www.pfizer.com.au](http://www.pfizer.com.au)

## 9. DATE OF FIRST APPROVAL

7 June 2018

## 10. DATE OF REVISION

04 July 2022

### Summary table of changes

Section changed	Summary of new information
Throughout	Minor editorial changes and spelling errors corrected
4.2	Heading "Method of Administration" added in alignment with TGA form for PI

4.2, 4.4 & 4.5	Information added regarding risk factors for QT interval prolongation
4.2, 4.3 & 4.4	Contraindication of children under 18 added
4.4	Warnings regarding haemolytic anaemia and positive Coombs' tests moved from Adverse Events information
4.4	Various other warnings and precautions added
4.5	Interaction warning added that neuroleptic medicinal products may have an antagonistic effect if used with apomorphine.
4.7	Warning added that apomorphine has varying degrees of impairment which influences the ability to drive and use machines
4.8	Various adverse events added
6.3	In use shelf life information repeated from Dose and Method of Administration section to Shelf Life section
6.7	Expression of chemical structure amended to match representation in Apomine Solution for Infusion PI