

FAMCICLOVIR MYLAN for cold sores

famciclovir

CONSUMER MEDICINE INFORMATION

What is in this leaflet

This leaflet answers some common questions about FAMCICLOVIR MYLAN.

It does not contain all the available information and does not take the place of talking to your doctor or pharmacist.

The information in this leaflet was last updated on the date listed on the final page.

Some more recent information on the medicine may be available.

You should ensure that you speak to your pharmacist or doctor to obtain the most up to date information on the medicine.

Those updates may contain important information about the medicine and its use of which you should be aware.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking FAMCICLOVIR MYLAN against the benefits it can provide.

If you have any concerns about taking this medicine, talk to your doctor or pharmacist.

Keep this leaflet with this medicine.

You may need to read it again.

What FAMCICLOVIR MYLAN is used for

FAMCICLOVIR MYLAN is an antiviral medicine used to treat recurrent outbreaks of cold sores in

adults and adolescents who have a normal immune system (the body system which fights against harmful bacteria, viruses and fungi).

Cold sores are an infection caused by a virus called herpes simplex type I (HSV-1). The infection is most commonly acquired as a baby or child from contact with parents or relatives, often from kissing.

Cold sores usually begin on or around the lips, mouth, and nose as small red bumps that turn into fluid-filled blisters. Cold sores can be tender and painful. Many people who get cold sores know when one is coming by a tingling, burning, itchy or painful sensation or redness in the area. This can happen very rapidly.

After redness and swelling develop, blisters form. The blisters may weep or burst and this can be painful. Then a shallow ulcer and yellow crust form as the cold sore dries. The crust eventually falls off, exposing new pink-coloured skin. Generally the sores heal without scarring.

After the initial infection has healed, the virus becomes dormant in nerve cells.

Cold sores can be unpredictable. The virus can become active again in the body, even after many years, resulting in recurrent outbreaks.

Even after many years, some people may experience recurring cold sores due to viral reactivation. Some common triggers to a cold sore may include:

- sun exposure
- stress

- fatigue
- menstrual periods
- fever
- illness
- dry chapped lips
- skin trauma
- a cold

Although FAMCICLOVIR MYLAN does not cure the viral infection, it helps to relieve the symptoms and shorten the duration of an outbreak.

The best results are obtained if the medicine is started as soon as possible after the onset of symptoms of a cold sore, such as tingling, itching or burning, or the appearance of the first signs, such as redness or swelling. This is when the virus is reproducing rapidly.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

FAMCICLOVIR MYLAN is only available with a doctor's prescription. It is not addictive.

This medicine is not recommended for use in infants, children or adolescents under 18 years of age.

Before taking FAMCICLOVIR MYLAN

When you must not take it

Do not take FAMCICLOVIR MYLAN if you have an allergy to:

- famciclovir, the active ingredient
- penciclovir, a related antiviral medicine
- any of the other ingredients of FAMCICLOVIR MYLAN listed at the end of this leaflet (see 'Product Description')

Some of the symptoms of an allergic reaction may include:

- shortness of breath, wheezing or difficulty breathing;
- swelling of the face, lips, tongue or other parts of the body;
- rash, itching or hives on the skin.

Do not take FAMCICLOVIR MYLAN after the expiry date printed on the pack, or if the packaging is torn or shows signs of tampering.

In that case, return it to your pharmacist.

Before you start to take it

Tell your doctor if you have:

- a problem with your body's immune system, which helps to fight off infections.
- a problem with your kidneys
- a problem with your liver

Your doctor may want to take extra precautions in that case.

Tell your doctor if you are pregnant, intend to become pregnant or if you are breast feeding.

FAMCICLOVIR MYLAN should not be used during pregnancy unless necessary. Your doctor will discuss with you the potential risks of taking FAMCICLOVIR MYLAN during pregnancy, and will also advise you if you should take FAMCICLOVIR MYLAN while breastfeeding, based on the benefits and risks of your particular situation.

Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.

Your doctor will want to know if you are prone to allergies.

If you experience an allergic reaction, inform your doctor or pharmacist immediately.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines and FAMCICLOVIR MYLAN may interfere with each other. These include:

- probenecid, a prescription medicine used to treat gout (a disease with painful, swollen joints caused by uric acid crystals) and to increase blood levels of penicillin-type antibiotics
- raloxifene, a medicine used to treat osteoporosis (a disease which causes bones to become less dense, gradually making them weaker, more brittle and likely to break)
- medicines that can affect your immune system
- medicines that can affect your kidneys

You may need to take different amounts of these medicines or you may need to take different medicines. Your doctor and pharmacist have more information.

If you have not told your doctor about any of these things, tell him/her before you start taking this medicine.

How to take FAMCICLOVIR MYLAN

Follow all directions given to you by your doctor and pharmacist carefully.

These instructions may differ from the information contained in this leaflet.

If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

How much to take

There are two ways to take FAMCICLOVIR MYLAN tablets to treat cold sores and your doctor will tell you which regimen is best for you:

- three 500 mg tablets taken together as a single dose
- or**
- two doses of 750 mg tablets taken 12 hours apart, for one day only.

However, your doctor may have prescribed a different dose.

For people whose immune system does not work as well as it should, the dose and duration of treatment may be increased.

For people who have kidney problems, your doctor may decide to give you a lower dose.

Ask your doctor or pharmacist if you are unsure of the correct dose for you.

They will tell you exactly how much to take.

Follow the instructions that they give you.

Do not change the dose yourself, without your doctor's advice, regardless of how well you may feel.

When to take it

Take FAMCICLOVIR MYLAN tablets as soon as possible after the first symptoms (e.g. tingling, itching or burning) or signs (e.g. redness or swelling) of a cold sore appear.

Do not take the tablets if a hard crust has already formed on the cold sore.

Keep the tablets for the next episode.

How to take it

Swallow the tablets whole with a full glass of water.

They may be taken with or without food.

How long to take it

A single dose of FAMCICLOVIR MYLAN for cold sores is all that is necessary for treating each episode of cold sores.

The dose may be repeated if cold sores recur. Each pack of FAMCICLOVIR MYLAN contains enough medicine for one dose.

If you take too much (Overdose)

Immediately telephone your doctor or the Australian Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at your nearest hospital, if you think that you or anyone else may have taken too much FAMCICLOVIR MYLAN. Show them your pack of tablets. Do this even if there are no signs of discomfort or poisoning. Keep the telephone numbers for these places handy.

Taking too much FAMCICLOVIR MYLAN may affect the kidneys. In people who already have kidney problems it may, rarely, lead to kidney failure if their dose is not correctly lowered.

While you are taking FAMCICLOVIR MYLAN

Things you must do

Tell your pharmacist or doctor if your cold sore symptoms do not improve within a few days, or if they become worse.

If you become pregnant while taking FAMCICLOVIR MYLAN, tell your doctor.

If you are about to be started on any new medicine, remind your doctor and pharmacist that you periodically take this medicine to treat recurring episodes of cold sores.

Tell any other doctor, dentist or pharmacist who treats you that you periodically take this medicine.

Things you must not do

Do not take less than the recommended dose of 3 tablets, unless advised by your doctor.

Do not give this medicine to anyone else even if their condition seems to be the same as yours.

Do not use this medicine to treat any other complaints unless your doctor tells you to.

Things to be careful of

If you are pregnant or breastfeeding, ask your doctor or pharmacist for advice before taking any medicine.

Be careful driving, operating machinery or doing jobs that require you to be alert until you know how FAMCICLOVIR MYLAN affects you.

This medicine can cause dizziness, sleepiness or confusion.

Things that may help your condition

Cold sores are contagious and the virus can be passed on from person to person through close physical contact or saliva, even when blisters are not present.

The risk is much higher when the cold sore is visible, as the virus can be shed, making it easy to infect other people.

Take the following precautions to avoid spreading the virus:

- Keep the areas affected by the virus as clean and dry as possible.
- Avoid touching or scratching the sore area as you may spread the virus on your fingers.
- Do not share any objects that have been in contact with a cold sore (e.g. drinking glasses, eating utensils, or towels).
- Avoid direct skin-to-skin contact of the area with other people (e.g.

kissing) until the cold sore has healed.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking FAMCICLOVIR MYLAN.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Do not be alarmed by these lists of possible side effects. You may not experience any of them. Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- headache
- dizziness
- nausea (feeling sick) or vomiting
- abdominal pain
- diarrhoea
- itching or an itchy rash (urticaria)

The above side effects are usually mild.

Tell your doctor as soon as possible if you notice any of the following:

- a rash elsewhere on the body, that is separate from the cold sore
- extreme sleepiness or confusion, usually in older people
- hallucinations (seeing or hearing things that are not really there)
- painful or swollen joints
- aching muscles or muscle tenderness or weakness that is not caused by exercise
- yellowing of the skin or eyes (signs of jaundice)
- palpitations (signs of abnormal heart beat)

The above side effects may need medical attention.

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if any of the following side effects happen:

- swelling below the surface of the skin (e.g. swelling around the face, eye, eyelid or throat)
- unexplained bruising, reddish or purplish patches on the skin or bleeding more easily than usual, as it may indicate that the number of platelets (a type of blood cell responsible for blood clotting) in your blood are reduced
- severe blistering of the skin or mucous membranes of the lips, eyes, mouth, nasal passages or genitals (signs of a serious skin reaction)
- purple patches, itching, burning of the skin (signs of inflamed blood vessels)

The above side effects are very rare.

Tell your doctor if you notice anything else that is making you feel unwell.

Other side effects not listed here or not yet known may happen in some people.

After taking FAMCICLOVIR MYLAN

Storage

Keep your medicine in the original container until it is time to take it.

Store your FAMCICLOVIR MYLAN tablets in a dry place where the temperature stays below 25°C.

Do not store your medicine in the bathroom or near a sink.

Do not leave the tablets in the car or on window sills.

Heat and dampness can destroy some medicines. FAMCICLOVIR MYLAN tablets will keep best if they are stored cool and dry.

Keep the medicine where children cannot reach it.

A locked cupboard at least one-and-a-half meters above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine you have left over.

Product description

What it looks like

FAMCICLOVIR MYLAN tablets are available in 500mg - white, oval shape, biconvex, film coated, debossed with "FAM 500" on one side and "G" on other side in blister packs of 3.

Ingredients

The active ingredient in FAMCICLOVIR MYLAN is famciclovir.

Each FAMCICLOVIR MYLAN 500 tablet contains 500 mg of famciclovir.

The tablets also contain the following inactive ingredients:

- hypolose
- sodium starch glycollate
- microcrystalline cellulose
- silicon dioxide
- magnesium stearate
- Opadry II white OY-L-28900

This product contains lactose.

Supplier

FAMCICLOVIR MYLAN is supplied in Australia by:

Alphapharm Pty Ltd

Level 1, 30 The Bond
30-34 Hickson Road
Millers Point NSW 2000
www.mylan.com.au

Australian registration numbers:

FAMCICLOVIR MYLAN 500 mg -
AUST R 157787

This leaflet was prepared on
13 May 2021.

FAMCICLOVIR MYLAN (cold sores) cmi\May21