

# Mersynofen<sup>®</sup>

**Active ingredients:** *ibuprofen and paracetamol*

---

## Consumer Medicine Information (CMI)

This leaflet provides important information about using Mersynofen. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Mersynofen.**

**Where to find information in this leaflet:**

- [1. Why am I using Mersynofen?](#)
- [2. What should I know before I use Mersynofen?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Mersynofen?](#)
- [5. What should I know while using Mersynofen?](#)
- [6. Things you should not do](#)
- [7. Are there any side effects?](#)
- [8. Product details](#)

## 1. Why am I using Mersynofen?

**Mersynofen contains the active ingredients ibuprofen and paracetamol.**

Ibuprofen belongs to a group of medicines called non-steroidal anti-inflammatory drugs (NSAIDs).

Paracetamol works to stop the pain messages from getting through to the brain.

Your doctor may have prescribed this medicine for another use.

**Mersynofen is an analgesic (pain reliever). It works to relieve acute (short term) pain and/or inflammation associated with headache, migraine headache, tension headache, sinus pain, toothache, dental procedures, backache, muscular aches and pains, period pain, sore throat, tennis elbow, rheumatic pain and arthritis, and the aches and pains associated with colds and flu.**

This medicine is not addictive.

## **2. What should I know before I use Mersynofen?**

### **Warnings**

#### **Do not use Mersynofen if:**

- you are allergic to paracetamol, ibuprofen, aspirin or any other medicine for pain relief, or any of the ingredients listed at the end of this leaflet  
Always check the ingredients to make sure you can use this medicine.
- you have a history of hypersensitivity reactions e.g. wheezing, chronic cough, swelling of the face or throat, red or itchy skin, skin rash, or a runny or stuffy

nose when using aspirin, or other non-steroidal anti-inflammatory drugs or analgesic products.

- you have liver or kidney disease
- you have pre-existing heart problems
- you have asthma
- you have a stomach ulcer or duodenal ulcer or if you have had either of these conditions or gastric bleeding or other gastrointestinal diseases in the past.
- you have recently vomited blood or material that looks like coffee grounds
- you have recently bled from the back passage (rectum), had black sticky bowel motions or bloody diarrhoea.
- you have had bleeding episodes which cannot be explained
- you are using other paracetamol-containing products
- you are severely dehydrated after having vomited, had diarrhoea or not enough to drink
- if the expiry date (EXP) printed on the pack or if the packaging is torn or shows signs of tampering

**Check with your doctor or pharmacist if you:**

- have a severe and extensive skin rash known as DRESS with following reactions such as fever, rash, enlargement of lymph nodes and/or facial swelling, kidney problems, blood disorder, inflammation of the heart, muscle weakness/pain
- have any other medical conditions, especially asthma, diabetes, liver or kidney disease or heart, gastrointestinal or autoimmune problems (where your

own immune system mistakenly attacks substances naturally occurring in your body).

- take any medicines for any other condition
- have recently had major surgery
- suffer from hayfever, nasal polyps or have chronic respiratory disorders. These may increase the risk of an allergic reaction occurring
- have a metabolic disorder
- have a blood platelet disorder
- consume large amounts of alcohol

**Tell your doctor or pharmacist if you have an infection.**

Mersynofen may hide signs of infections such as fever and pain.

It is therefore possible that Mersynofen may delay appropriate treatment of infection, which may lead to an increased risk of complications.

This has been observed in patients with serious lung infections (also called pneumonia) caused by bacteria and bacterial skin infections related to chickenpox. If you take this medicine while you have an infection and your symptoms of the infection persist or worsen, consult a doctor without delay.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

## **Pregnancy and breastfeeding**

**Do not take Mersynofen if you are pregnant or intend to become pregnant.**

Like most medicines of this kind, Mersynofen is not recommended to be used during pregnancy.

Use of Mersynofen may cause the baby to have kidney problems leading to a low amount of amniotic fluid inside the uterus and in some cases kidney problems in the newborn.

**Check with your doctor or pharmacist about taking Mersynofen if you are breast-feeding or planning to breast-feed.**

Small amounts of ibuprofen and paracetamol may pass into the breast milk

**If you have not told your doctor or pharmacist about any of the above, tell them before start taking Mersynofen.**

## **Use in the elderly**

**Do not take if you are aged 65 years or older.**

## **Use in children**

**Do not give Mersynofen to children aged under 12 years.**

**Do not give Mersynofen to dehydrated adolescents.**

### **3. What if I am taking other medicines?**

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

**Some medicines and Mersynofen may interfere with each other. These include:**

- other paracetamol containing products, analgesics, other medicines for pain relief
- warfarin, a medicine used to prevent blood clots
- antiplatelet agents and selective serotonin reuptake inhibitors (SSRIs)
- lithium, a medicine used to treat mood swings and some types of depression
- medicines used to lower blood pressure such as diuretics, ACE inhibitors, beta-receptor blockers and angiotensin-II antagonists
- medicines that contain probenecid or sulfinpyrazone
- salicylates or non-steroidal anti-inflammatory drugs (NSAIDS), such as aspirin
- digoxin, a medicine used to treat irregular heart rhythms
- medicines used to lower cholesterol such as cholestyramine
- methotrexate, a medicine used to treat arthritis and some types of cancer
- medicines used to treat heart failure

- corticosteroids such as prednisone, prednisolone and cortisone, which reduces the activity of your immune system
- zidovudine, a medicine used to treat HIV infection
- aspirin, salicylates and other non-steroidal anti-inflammatory drugs (NSAIDs)
- medicines used to treat diabetes
- metoclopramide, a medicine used to control nausea and vomiting
- medicines used to treat epilepsy or fits (seizures)
- chloramphenicol an antibiotic used to treat ear and eye infections
- ciclosporin or tacrolimus, medicines used to treat immune system conditions
- alcohol
- CYP2C9 inhibitors such as voriconazole and fluconazole
- uricosurics which may delay the excretion of ibuprofen

These medicines may be affected by Mersynofen or may affect how well Mersynofen works. You may need different amounts of your medicines, or you may need to take different medicines.

**If you have not told your doctor or pharmacist about any of these things, tell him/her before you take any Mersynofen.**

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Mersynofen.**

## 4. How do I use Mersynofen?

### How much to take

- The label on your pack of Mersynofen will tell you how to take your medicine and how often.

If you do not understand the instructions on the pack, ask your doctor or pharmacist for help.

- **The usual dose of Mersynofen is:**

Adults under 65 and children from 12 years:

- 1 tablet.
- This dosage may be repeated every 8 hours if necessary.

**You should not take more than 3 tablets in 24 hours.**

Do not take more than the recommended dose.

Not recommended for adults 65 years of age and over or children under 12 years of age.

### How to take Mersynofen

- For oral use only
- Swallow tablets with a little water or other liquid.
- It is recommended that patients with sensitive stomachs take Mersynofen with food.
- If taken shortly after eating it may take a little longer for Mersynofen to work. Do not take more Mersynofen than recommended until it is time to take the next dose.



## **How long to take it**

**Only take Mersynofen when required for pain.**

Take the lowest dose that helps with your pain for the shortest time necessary to relieve your symptoms.

If you have an infection, consult a doctor without delay if symptoms (such as fever and pain) persist or worsen.

### **Adults:**

Do not take Mersynofen for more than 3 days at a time.

### **Adolescents (12 to 17 years):**

Do not take Mersynofen for more than 2 days at a time.

**If your symptoms persist, worsen or new symptoms develop, talk to your pharmacist or doctor.**

Prolonged use of any type of pain relievers for headaches can make it worse. If this is experienced or suspected, medical advice should be obtained or discontinued.

The habit of taking medicines for pain relief particularly in combination with other pain-relievers may lead to permanent kidney problems including kidney failure therefore it should be avoided.

## **If you take too much Mersynofen**

If you think that you have taken too much Mersynofen, you may need urgent medical attention.

**You should immediately:**

- phone the Poisons Information Centre **(by calling 13 11 26)**, or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

**You should do this even if there are no signs of discomfort or poisoning.**

## **5. What should I know while using Mersynofen?**

### **Things you should do**

**If you are about to be started on any new medicine, tell your doctor and pharmacist that you are taking Mersynofen.**

**Talk to your doctor or pharmacist if your symptoms don't improve.**

Your pharmacist or doctor will assess your condition and decide if you should continue to take the medicine.

**If you become pregnant while you are using this medicine, tell your doctor or pharmacist immediately.**

**If you develop a skin rash, mucosal wounds or any other sign of allergy while you are using this medicine, stop taking the product and tell your doctor or pharmacist immediately.**

## **Things you should not do**

- **Do not take for more than 3 days at a time (2 days for adolescents 12 to 17 years) unless your doctor or pharmacist tells you to.**
- **Do not take more than the recommended dose unless your doctor or pharmacist tells you to.**

## **Things to be aware of**

Taking this medicine may increase the risk of you getting unwanted effects, such as stomach or heart problems.

**Do not take Mersynofen to treat any other complaints unless your pharmacist or doctor tells you to.**

**Do not give your medicine to anyone else, even if they have the same condition as you.**

## **Driving or using machines**

**Be careful driving or operating machinery until you know how this medicine affects you.**

This medicine may cause tiredness or dizziness in some people.

## **Drinking alcohol**

**Only drink small quantities of alcohol (beer, wine or spirits) while taking paracetamol.**

Drinking large quantities of alcohol while taking paracetamol may increase the risk of liver side effects.

## **Looking after your medicine**

- Store below 30°C.

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on windowsills.

**Keep it where young children cannot reach it.**

## **Getting rid of any unwanted medicine**

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

## **6. Are there any side effects?**

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

## Less serious side effects

| Less serious side effects   | What to do   |
|---|--|
| <p data-bbox="204 432 762 477"><b>Gastrointestinal related:</b></p> <ul data-bbox="204 510 678 1104" style="list-style-type: none"><li data-bbox="204 510 411 555">● nausea</li><li data-bbox="204 577 481 622">● indigestion</li><li data-bbox="204 645 470 689">● heart burn</li><li data-bbox="204 712 539 757">● stomach pain</li><li data-bbox="204 779 454 824">● diarrhoea</li><li data-bbox="204 846 577 891">● loss of appetite</li><li data-bbox="204 913 678 958">● wind or constipation</li><li data-bbox="204 981 577 1025">● <b>Eye Disorders</b></li><li data-bbox="204 1048 646 1093">● Visual disturbance</li></ul> <p data-bbox="204 1126 670 1227"><b>Head and neurology related:</b></p> <ul data-bbox="204 1261 523 1518" style="list-style-type: none"><li data-bbox="204 1261 450 1305">● dizziness</li><li data-bbox="204 1328 491 1373">● drowsiness</li><li data-bbox="204 1395 466 1440">● headache</li><li data-bbox="204 1462 523 1507">● nervousness</li></ul> <p data-bbox="204 1541 555 1597"><b>Allergy related:</b></p> <ul data-bbox="204 1630 497 1742" style="list-style-type: none"><li data-bbox="204 1630 497 1675">● skin rashes</li><li data-bbox="204 1697 443 1742">● sweating</li></ul> <p data-bbox="204 1765 577 1865"><b>Immune System Disorders</b></p> <ul data-bbox="204 1899 646 2011" style="list-style-type: none"><li data-bbox="204 1899 646 2011">● Hypersensitivity reactions with skin</li></ul> | <p data-bbox="817 432 1375 645"><b>Speak to your doctor if you have any of these less serious side effects and they worry you.</b></p> |

| <b>Less serious side effects</b>   | <b>What to do</b> |
|--|-------------------|
| <p>rashes and itching, as well as asthma attacks (possibly with drop in blood pressure)</p> <p>Pregnancy related</p> <ul style="list-style-type: none"> <li>● low amniotic fluid inside uterus (oligohydramnios)</li> <li>● kidney problems in newborn baby</li> </ul> |                   |

## **Serious side effects**

| <b>Serious side effects</b>  | <b>What to do</b>  |
|--|--|
| <p><b>Allergy related:</b></p> <ul style="list-style-type: none"> <li>● shortness of breath</li> <li>● wheezing or difficulty breathing</li> <li>● swelling of the face, lips, tongue or other parts of the body</li> <li>● rash, itching or hives on the skin</li> <li>● flushing of the face</li> </ul> <p><b>Heart related:</b></p> <ul style="list-style-type: none"> <li>● fluid retention</li> </ul> | <p><b>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</b></p> |

| <b>Serious side effects</b>  | <b>What to do</b> |
|--|-------------------|
| <p><b>Breathing related:</b></p> <ul style="list-style-type: none"> <li>● shortness of breath</li> <li>● wheezing or difficulty breathing</li> </ul> <p><b>Skin related:</b></p> <ul style="list-style-type: none"> <li>● you are getting sunburnt more quickly than usual.</li> <li>● drug induced reddish, scaly, pus-filled bumps</li> </ul> <p><b>Gastrointestinal related:</b></p> <ul style="list-style-type: none"> <li>● vomiting blood or bleeding from the back passage</li> </ul> |                   |

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

### **Reporting side effects**

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at

[www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

## **7. Product details**

This medicine is only available from your pharmacist.

### **What Mersynofen contains**

|   |  |
|---|--|
| <b>Active ingredients<br/>(main ingredient)</b>     | Ibuprofen and paracetamol.   |
| <b>Other ingredients<br/>(inactive ingredients)</b> | Maize starch, magnesium stearate , microcrystalline cellulose, pregelatinised maize starch, croscarmellose sodium, sodium lauryl sulfate, purified talc, opadry QX Quick and Flexible Film Coating System 321A180025 White, and opadry fx special effects film coating system 63F97546 Silver. |



**Do not take this medicine if you are allergic to any of these ingredients.**

## **What Mersynofen looks like**

Mersynofen is a white to off white, capsule shaped, film coated tablet with a break bar on one side (AUST R 321566).

Mersynofen is available in a range of sizes up to 30 tablets.

## **Who distributes Mersynofen**

Sanofi Consumer Healthcare,

87 Yarraman Place

Virginia, QLD 4014, Australia

Toll Free Number (medical information): 1800 818 806

Email: [medinfo.australia@sanofi.com](mailto:medinfo.australia@sanofi.com)

This leaflet was prepared in October 2021.

mersynofen-cmiv4-ibuccdsv4paraccdsv3-21oct21