

# CILICAINE VK

phenoxymethylpenicillin (as potassium)

## Consumer Medicine Information

### What is in this leaflet

This leaflet answers some common questions about CILICAINE VK. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of using CILICAINE VK against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

### What CILICAINE VK is used for

This medicine contains the active ingredient phenoxymethylpenicillin. It is a type of antibiotic that belongs to the group of medicines called penicillins.

CILICAINE VK is used to treat infections in different parts of the body caused by bacteria. It works by killing the bacteria that are causing the infection.

CILICAINE VK will not work against infections caused by viruses such as colds or the flu.

**Ask your doctor if you have any questions about why this medicine has been prescribed for you.**

Your doctor may have prescribed CILICAINE VK for another purpose.

This medicine is available only with a doctor's prescription.

### Before you take CILICAINE VK

#### **When you must not take it**

**Do not take CILICAINE VK if you have an allergy to:**

- any medicine containing phenoxymethylpenicillin
- any other penicillin medicines or cephalosporins
- any of the ingredients listed at the end of this leaflet

**If you have an allergic reaction to cephalosporin you may have an increased chance of being allergic to CILICAINE VK.**

Some symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

**Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.**

If it has expired or is damaged, return to your pharmacist for disposal.

**If you are not sure whether you should start taking this medicine, talk to your doctor.**

#### **Before you start to take it**

**Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

**Tell your doctor if you have or have had any of the following medical conditions:**

- diarrhoea after taking antibiotics previously
- bleeding disorders
- kidney problems
- liver problems
- any other health problem

**Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.**

Your doctor can discuss with you the risks and benefits involved.

The active ingredient in CILICAINE VK passes into breast milk and there is a possibility that your baby may be affected.

**If you have not told your doctor about any of the above, tell them before you start taking CILICAINE VK.**

#### **Taking other medicines**

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop**

Some medicines and CILICAINE VK may interfere with each other. These include:

- probenecid, a medicine used to treat gout
- antacids, medicines used to reduce heartburn or indigestion

- oral contraceptive pills, medicines used for contraception

You may need to use other birth control methods while you are taking CILICAINE VK.

- aminoglycosides, medicines used to treat some bacterial infections
- methotrexate, a medicine used to treat rheumatoid arthritis, some cancers and inflammatory conditions

These medicines may be affected by CILICAINE VK or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

---

## How to take CILICAINE VK

---

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the box, ask your doctor or pharmacist for help.**

### **How much to take**

The usual dose is:

#### **Adults**

250 mg to 500 mg every four to six hours.

#### **To prevent recurrence of rheumatic fever**

250 mg twice a day for as long as your doctor tells you to.

The dose you take will depend on the type of treatment you are to receive. Your doctor will decide your dose and how long you take CILICAINE VK for.

### **How to take it**

Swallow the capsules whole with a glass of water.

### **When to take it**

Take your medicine 1 hour before food.

### **How long to take it**

Continue taking your medicine until you finish the pack or for as long as your doctor tells you.

It is important to keep taking your medicine even if you feel well.

### **If you forget to take it**

If it is less than 4 hours before your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose you missed.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

### **If you take too much (overdose)**

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much CILICAINE VK. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

---

## While you are using CILICAINE VK

---

### **Things you must do**

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking CILICAINE VK.

Tell any other doctors, dentists and pharmacists who treat you that you are taking this medicine.

If you become pregnant while taking this medicine, tell your doctor immediately.

If your symptoms do not improve within a few days, or if they become worse, tell your doctor immediately.

If you get diarrhoea, nausea or vomiting, tell your doctor or pharmacist immediately. Do this even if it occurs several weeks after you have stopped taking CILICAINE VK.

Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care.

If you are about to have any blood or urine tests, tell your doctor that you are taking this medicine.

It may interfere with the results of some tests.

### **Things you must not do**

Do not take any medicine for diarrhoea without first checking with your doctor or pharmacist.

Do not take CILICAINE VK to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking CILICAINE VK without checking with your doctor.

Your infection and symptoms may not clear completely.

### **Things to be careful of**

**Tell your doctor if you get a sore mouth or tongue while taking or after you have stopped taking CILICAINE VK.**

This could be due to a fungal infection called thrush. This may also result in fever and irritation in your stomach.

**Be careful driving or operating machinery until you know how CILICAINE VK affects you.**

---

### **Side effects**

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking CILICAINE VK.**

This medicine helps most people but it may have some unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

**Do not be alarmed by the following lists of side effects. You may not experience any of them.**

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor or pharmacist if you notice any of the following and they worry you:**

- skin rash
- nausea or vomiting
- black hairy tongue
- diarrhoea
- upset stomach
- difficulty in breathing

**Tell your doctor or pharmacist if you notice anything that is making you feel unwell.**

Other side effects not listed above may also occur in some people.

---

## **After using CILICAINE VK**

---

### **Storage**

**Keep your capsules in the pack until it is time to take them.**

If you take the capsules out of the pack they may not keep well.

**Keep your capsules in a cool dry place where the temperature stays below 25°C.**

**Do not store CILICAINE VK or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.**

Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### **Disposal**

**If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.**

---

## **Product description**

---

### **What it looks like**

CILICAINE VK 250 mg and 500 mg capsules are maroon in colour.

### **Ingredients**

CILICAINE VK contains 250 mg or 500 mg of phenoxymethylpenicillin as the active ingredient.

The capsules also contain the following inactive ingredients:

- gelatin
- titanium dioxide
- iron oxide red
- magnesium stearate

### **Distributor**

CILICAINE VK is distributed in Australia by:

**Alphapharm Pty Ltd**

Level 1, 30 The Bond

30-34 Hickson Road

Millers Point NSW 2000

www.mylan.com.au

This leaflet was prepared in May 2020.

250 mg capsules: AUST R 55676

500 mg capsules: AUST R 55675

**cilicaine-vk\_cmi\May20/00**