

APO-Candesartan HCTZ Tablets

Contains the active ingredient candesartan cilexetil and hydrochlorothiazide

Consumer Medicine Information

For a copy of a large print leaflet, Ph: 1800 195 055

What is in this leaflet

Read this leaflet carefully before taking your medicine.

This leaflet answers some common questions about candesartan cilexetil and hydrochlorothiazide (HCTZ). It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

The information in this leaflet was last updated on the date listed on the last page. More recent information on this medicine may be available.

Ask your doctor or pharmacist:

- if there is anything you do not understand in this leaflet,
- if you are worried about taking your medicine, or
- to obtain the most up-to-date information.

All medicines have risks and benefits. Your doctor has weighed the risks of you using this medicine against the benefits they expect it will have for you.

Pharmaceutical companies cannot give you medical advice or an individual diagnosis.

Keep this leaflet with your medicine. You may want to read it again.

What this medicine is used for

The name of your medicine is APO-Candesartan HCTZ tablets. It contains the active ingredients candesartan cilexetil and hydrochlorothiazide.

It is used to treat:

- high blood pressure, also called hypertension.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed this medicine for another reason.

This medicine is available only with a doctor's prescription.

How it works

Candesartan cilexetil is a type of medicine called an angiotensin II receptor inhibitor (or antagonist). It mainly works by causing relaxation of blood vessels.

Hydrochlorothiazide is a type of medicine called a diuretic. It works by reducing the amount of excess fluid in the body.

Using these two medicines together will lower your blood pressure more effectively than using either one on its own.

There is no evidence that this medicine is addictive.

Use in children

This medicine should not be used in children. The safety and effectiveness of this medicine in children under 18 years of age have not been established.

Before you use this medicine

When you must not use it

Do not use this medicine if:

- **You have or have had any of the following:**
- severe kidney disease
- gout
- severe liver disease and/or cholestasis.

- taking blood pressure medicine containing aliskiren, especially if you have diabetes mellitus or kidney problems.
- **You are pregnant or planning to become pregnant.**

This medicine may affect your developing baby if you take it during pregnancy.

- **You are breast-feeding or planning to breast-feed.**

This medicine may pass into human breast milk.

- **You are hypersensitive to, or have had an allergic reaction to, candesartan cilexetil, hydrochlorothiazide or any of the ingredients listed at the end of this leaflet.**
- **You are hypersensitive to, or have had an allergic reaction to, any sulphur drugs (sulphonamides) such as antibiotics or some medicines used to treat diabetes.**

Symptoms of an allergic reaction may include cough, shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue, throat or other parts of the body; rash, itching or hives on the skin; fainting or hayfever-like symptoms.

If you think you are having an allergic reaction, do not take any more of the medicine and contact your doctor immediately or go to the Accident and Emergency Department at the nearest hospital.

- **The expiry date (EXP) printed on the pack has passed.**

- The packaging is torn, shows signs of tampering or it does not look quite right.

Before you start to take it

Before you start taking this medicine, tell your doctor if:

1. You have allergies to:
 - any other medicines
 - any other substances, such as foods, preservatives or dyes.
2. You have or have had any medical conditions, especially the following:
 - kidney problems
 - liver problems
 - heart problems
 - diabetes
 - recent excessive vomiting or diarrhoea
 - Systemic Lupus Erythematosus (SLE), a disease affecting the skin, joints and kidneys
 - a salt restricted diet
 - a condition called primary hyperaldosteronism
 - a past operation known as sympathectomy

- you are hypersensitive to, or have had an allergic reaction to, penicillin.
- 3. You are currently pregnant or you plan to become pregnant. Do not take this medicine whilst pregnant.
- 4. You are currently breast-feeding or you plan to breast-feed. Do not take this medicine whilst breast-feeding.
- 5. You have recently been vaccinated or plan to get a vaccination.
- 6. You are planning to have surgery or an anaesthetic.
- 7. You are currently receiving or are planning to receive dental treatment.
- 8. You are taking or are planning to take any other medicines. This includes vitamins and supplements that are available from your pharmacy, supermarket or health food shop.

Some medicines may interact with Candesartan HCTZ.. These include:

- other medicines that lower blood pressure such as ACE inhibitors, angiotensin II receptor blockers and aliskiren, especially if you have diabetes-related kidney problems
- medicines used to raise blood pressure.
- other medicines associated with potassium loss, such as other diuretics and laxatives.

- medicines containing potassium, including salt substitutes.
- cardiac glycosides, medicines used to treat heart failure and irregular heartbeat, such as Digoxin.
- other medicines used to treat irregular heartbeat.
- non-steroidal anti-inflammatory drugs (NSAIDs), medicines used to relieve pain, swelling and other symptoms of inflammation, including arthritis.
- colestipol and cholestyramine, medicines used to treat high blood cholesterol levels.
- lithium, a medicine used to treat mood swings and some types of depression.
- alcohol.
- strong pain killers such as codeine, morphine, dextropropoxyphene.
- barbiturates, used to treat epilepsy, such as phenobarbitone.
- medicines like insulin that are used to treat diabetes.
- calcium supplements, or medicines containing calcium.
- Vitamin D supplements
- corticosteroids such as prednisone, cortisone, dexamethasone.

- non-depolarizing muscle relaxants such as tubocurarine.
- iodinated contrast media, a radiographic dye used for imaging techniques such as CT scans.
- amantadine, an antiviral and an antiparkinsonian medicine.
- anticholinergic agents, medicines that act on the nervous system, such as atropine.
- cytotoxic medicines, such as medicines used for chemotherapy.
- cyclosporine, an immunosuppressant used to prevent organ transplant rejection.

If you are taking any of these you may need a different dose or you may need to take different medicines.

Other medicines not listed above may also interact with candesartan HCTZ.

How to take this medicine

Follow carefully all directions given to you by your doctor carefully.

Their instructions may be different to the information in this leaflet.

If you are not sure how to take this medicine, ask your doctor or pharmacist for help.

How much to take

Your doctor or pharmacist will tell you how much of this medicine you should take. This will depend on your condition and whether you are taking any other medicines.

Do not stop taking your medicine or change your dosage without first checking with your doctor.

The usual dose is one tablet once daily.

How to take it

Swallow the tablets whole with a glass of water.

When to take it

Take this medicine at the same time each day. Taking it at the same time each day will have the best effect and will also help you remember when to take it.

It does not matter if you take it before, with or after food.

How long to take it

Continue taking this medicine for as long as your doctor tells you.

This medicine controls your condition, but does not cure it. It is important to keep taking your medicine even if you feel well.

Make sure you have enough to last over weekends and holidays.

If you forget to take it

If you forget to take a dose, take it as soon as you remember, as long as it is at least 12 hours before your next dose is due.

Then go back to taking it as you would normally.

If it is less than 12 hours to your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for missed doses. This may increase the chance of you experiencing side effects.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints to help you remember.

If you take too much (overdose)

If you think that you or anyone else may have taken too much of this medicine, immediately telephone your doctor or the Poisons Information Centre (Tel: 13 11 26 in Australia) for advice. Alternatively go to the Accident and Emergency Department at your nearest hospital.

Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

If you take too many candesartan HCTZ tablets, you may get a headache, feel sick (nausea), dizzy, thirsty and very tired.

While you are taking this medicine

Things you must do

Tell your doctor that you are taking this medicine if:

- you are about to be started on any new medicine
- you are pregnant or are planning to become pregnant
- you are breast-feeding or are planning to breast-feed
- you are about to have any blood tests
- you are going to have surgery or an anaesthetic or are going into hospital
- you experience excessive vomiting or diarrhoea.
- You may lose too much water and your blood pressure may become too low.

Your doctor may occasionally do tests to make sure the medicine is working and to prevent side effects. Go to your doctor regularly for a check-up.

Tell any other doctors, dentists and pharmacists who are treating you that you take this medicine.

Things you must not do

Do not:

- Give this medicine to anyone else, even if their symptoms seem similar to yours.
- Take your medicine to treat any other condition unless your doctor tells you to.
- Stop taking your medicine, or change the dosage, without first checking with your doctor.

Things to be careful of

Be careful driving or operating machinery until you know how candesartan HCTZ affects you.

You may feel dizzy when you start taking candesartan HCTZ due to the drop in your blood pressure.

Move slowly when getting out of bed or standing up if you feel faint, dizzy or light-headed.

Drink plenty of water while you are using this medicine, especially if you sweat a lot.

Please talk to your doctor or pharmacist about these possibilities if you think they may be a problem for you.

Possible side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking this medicine or if you have any questions or concerns.

Do not be alarmed by the following lists of side effects. You may not experience any of them. All medicines can have

side effects. Sometimes they are serious but most of the time they are not.

Tell your doctor if you notice any of the following:

- headache or dizziness
- flu-like symptoms or infections
- chest, throat or sinus infections
- feeling sick (nausea)
- back pain
- urinary tract infection
- rapid heartbeats
- feeling tired
- stomach ache.

Tell your doctor as soon as possible if you notice any of the following:

- aching muscles, tenderness or weakness in the muscle.

The above list includes serious side effects and are usually rare. They may require medical attention.

If you experience any of the following, stop taking your medicine and contact your doctor immediately or go to Accident and Emergency at your nearest hospital:

- easy bruising or bleeding more easily than normal
- feeling extremely tired
- yellowing of the skin and eyes
- signs of frequent infections such as fever, severe chills, sore throat or mouth ulcers
- worsening of the kidney function including passing little or no urine, drowsiness, nausea, vomiting, breathlessness, loss of appetite and weakness (especially in patients with existing kidney problems or heart failure)
- changes in your potassium, sodium and red or white blood cell levels may occur. Such changes are usually detected by a blood test
- symptoms that may indicate high potassium levels in the blood include nausea, diarrhoea, muscle weakness and changes in heart rhythm
- blurred vision or eye pain.

These are very serious side effects and are usually very rare. You may need urgent medical attention or hospitalisation.

Other side effects not listed above may above may occur in some patients. Tell your doctor if you notice any other side effects.

Allergic reactions

If you think you are having an allergic reaction to candesartan HCTZ, do not take any more of this medicine

and tell your doctor immediately or go to the Accident and Emergency department at your nearest hospital.

Symptoms of an allergic reaction may include some or all of the following:

- cough, shortness of breath, wheezing or difficulty breathing
- swelling of the face, lips, tongue, or other parts of the body
- rash, itching or hives on the skin
- fainting
- hayfever-like symptoms.

Storage and disposal

Storage

Keep your tablets in the blister pack until it is time to take them.

If you take candesartan out of the blister pack it will not keep well.

Keep your tablets in a cool dry place where the temperature will stay below 25°C.

Do not store your medicine, or any other medicine, in the bathroom or near a sink. Do not leave it on a window sill or in the car. Heat and dampness can destroy some medicines.

Keep this medicine where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor or pharmacist tells you to stop taking this medicine or it has passed its expiry date, your pharmacist can dispose of the remaining medicine safely.

Product description

What APO-Candesartan HCTZ looks like

APO-Candesartan HCTZ 16/12.5 mg tablets: light pink, oval shape, biconvex, uncoated, mottled tablet debossed with 'L3' and '02' on either side of breakline on one side and break line on other side APO-Candesartan HCTZ 32/12.5 mg tablets: light yellow, oval, biconvex, uncoated, mottled tablet debossed with 'L3' & '04' on either side of breakline on one side and break line on other side APO-Candesartan HCTZ 32/25 mg tablets: light pink, oval, biconvex, uncoated, mottled tablet debossed with 'L3' & '04' on either side of break line on one side & break line on one other side Ingredients

APO-Candesartan HCTZ Tablets 16/12.5 mg contains 16 mg candesartan cilexetil and 12.5 mg of hydrochlorothiazide.

APO-Candesartan HCTZ Tablets 32/12.5 mg contains 32 mg candesartan cilexetil and 12.5 mg of hydrochlorothiazide.

APO-Candesartan HCTZ Tablets 32/25 mg contains 32 mg candesartan cilexetil and 25 mg of hydrochlorothiazide.

APO-Candesartan HCTZ is available in:

Blister packs of 7 and 30 tablets.

Not all strengths, and/or pack sizes may be available.

Ingredients

Each tablet contains 16 mg or 32mg of candesartan and 12.5mg or 25mg of HCTZ as the active ingredient.

It also contain the following inactive ingredients:

- lactose
- carmellose calcium
- starch - maize
- macrogol 8000
- hydroxypropylcellulose
- magnesium stearate
- Pigment Blend PB-24880 Pink (16/12.5 mg and 32/25 tablet only)
- (iron oxide yellow) (32/12.5 mg tablet only).

This medicine is gluten-free, sucrose-free, tartrazine-free and free of other azo dyes.

Australian Registration Numbers

APO-Candesartan HCTZ 16/12.5 mg blister pack: AUST R 210565

APO-Candesartan HCTZ 32/12.5 mg blister pack: AUST R 210566

APO-Candesartan HCTZ 32/25 mg blister pack: AUST R 210567

Sponsor

Apotex Pty Ltd

16 Giffnock Avenue

Macquarie Park, NSW 2113

Australia

Apotex Pty Ltd is the licensee of the registered trademarks APO and APOTEX from the registered proprietor, Apotex Inc.

This leaflet was prepared in November 2016.