

Mersyndol® DayStrength

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using Mersyndol DayStrength?

Mersyndol DayStrength contains the active ingredients paracetamol and codeine phosphate hemihydrate. Mersyndol DayStrength is used to relieve acute moderate pain and fever.

For more information, see Section [1. Why am I using Mersyndol DayStrength?](#) in the full CMI.

2. What should I know before I use Mersyndol DayStrength?

Do not use if you have ever had an allergic reaction to Mersyndol DayStrength or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use Mersyndol DayStrength?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Mersyndol DayStrength and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use Mersyndol DayStrength?

Adults and children 12 years or over:

- One or two tablets every 3 to 4 hours as needed for relief. Do not take more than 8 caplets in 24 hour period
- More instructions can be found in Section [4. How do I use Mersyndol DayStrength?](#) in the full CMI.

5. What should I know while using Mersyndol DayStrength?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using Mersyndol DayStrength.• Tell your doctor or pharmacist if you are taking any other medicines that you use to help you relax, anything that contains alcohol (like cough syrup) or other medicines that treat pain.
Things you should not do	<ul style="list-style-type: none">• Do not take more than the recommended dose unless your doctor tells you to.• Do not give this medicine to children under 12 years of age.• Do not take Mersyndol DayStrength during the third trimester of pregnancy or if you are in labour, especially if the baby is expected to be premature.
Driving or using machines	<ul style="list-style-type: none">• Mersyndol DayStrength may cause drowsiness in some people. If this happens, do not drive or operate machinery.
Drinking alcohol	<ul style="list-style-type: none">• Do not drink alcohol while taking Mersyndol DayStrength as it may interfere with the effects of the medicine and may increase the risk of liver side effects.
Looking after your medicine	<ul style="list-style-type: none">• Store below 30° C.• Store in a cool, dry place away from young children.

For more information, see Section [5. What should I know while using Mersyndol DayStrength?](#) in the full CMI.

6. Are there any side effects?

Mersyndol DayStrength may be habit forming if taken frequently or over long periods. Tell your doctor or pharmacist if you notice any of the following and they worry you: nausea or vomiting; drowsiness or dizziness; constipation; stomach pain; skin rashes; sweating. For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

Mersyndol DayStrength®

Active ingredient(s): *paracetamol and codeine phosphate hemihydrate*

Consumer Medicine Information (CMI)

This leaflet provides important information about using Mersyndol DayStrength. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Mersyndol DayStrength.**

Where to find information in this leaflet:

- [1. Why am I using Mersyndol DayStrength?](#)
- [2. What should I know before I use Mersyndol DayStrength?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Mersyndol DayStrength?](#)
- [5. What should I know while using Mersyndol DayStrength?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using Mersyndol DayStrength?

Mersyndol DayStrength is a type of analgesic intended for short term use to relieve acute moderate pain and fever.

Mersyndol DayStrength contains the active ingredients paracetamol and codeine phosphate hemihydrate.

Codeine phosphate hemihydrate belongs to a group of medicines called analgesics. It is an opioid analgesic and it acts by blocking pain and your emotional response to pain.

Paracetamol works to stop the pain messages from getting through to the brain. It also acts in the brain to reduce fever.

Paracetamol and codeine work together to stop the pain messages from getting through to the brain.

Mersyndol DayStrength is for the short term relief of acute moderate pain and fever.

2. What should I know before I use Mersyndol DayStrength?

Warnings

Do not use Mersyndol DayStrength if:

- you are allergic to paracetamol or codeine phosphate hemihydrate, or any of the ingredients listed at the end of this leaflet.

Always check the ingredients to make sure you can use this medicine.

- You are allergic to aspirin or anti-inflammatory medications
- you have acute breathing difficulties such as bronchitis, unstable asthma or emphysema (serious lung disease).
- you have a Glucose-6-phosphate dehydrogenase deficiency (an enzyme deficiency)
- you are a ultra-rapid metaboliser of CYP 2D6 (a fast metaboliser of codeine by the CYP 2D6 enzyme)
- you have had your tonsils or adenoids removed and are aged between 12 and 18 years of age
- you have severe liver or kidney disease
- you are suffering from diarrhea caused by poisoning or antibiotics

- you have chronic constipation

Do not take codeine if you have alcohol dependence.

Do not give Mersyndol Day Strength to children under 12 years of age.

Do not take Mersyndol Day Strength during the third trimester of pregnancy.

Do not take it during labour, especially if the baby is premature.

Do not take it if you are breastfeeding or planning to breastfeed.

Do not take this medicine after the expiry date (EXP) printed on the pack.

If you take it after the expiry has passed, it may not work as well.

Do not take this medicine if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

Check with your doctor if you:

- plan to have surgery
- have difficulty breathing, experience wheezing, chronic cough, asthma or other chronic breathing conditions.
- have any form of problems with breathing as a result of emphysema, kyphoscoliosis or obesity
- have a known analgesic intolerance
- have or have had heart, liver or kidney problems
- underactive thyroid
- have a history of drug dependence, including alcohol dependence
- experience chronic alcohol use including recent cessation of alcohol intake
- have a pre-existing opioid dependence
- have low glutathione reserves
- have Gilbert's syndrome
- have had recent surgery of the stomach or intestine
- have prostate problems
- have multiple sclerosis
- have low blood pressure
- have had a head injury or trauma
- have urinary, bowel or gallbladder conditions
- have chronic constipation
- have problems with the adrenal glands
- have myasthenia gravis, a muscle disorder
- have convulsions, fits or seizures

Tell your doctor or pharmacist about anything listed above before you start taking Mersyndol DayStrength.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed. Mersyndol DayStrength passes into breast milk and there is a possibility your baby may be affected.

Do not take Mersyndol DayStrength during the third trimester of pregnancy.

Do not take Mersyndol DayStrength during labour, especially if the baby is premature.

This may produce withdrawal effects in the newborn baby.

If you are over 65 years of age

Talk to your doctor or pharmacist about how much to take.

Elderly patients are more likely to have less effective kidney function due to age. This may increase the risk of side effects.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with the absorption of Mersyndol DayStrength. These include:

- medicines used to help relax, sleep or relieve anxiety, such as barbiturates and sedatives
- benzodiazepines (medicines used as sedatives or to treat anxiety)
- medicines containing alcohol (ethanol), e.g. some cough syrups
- antihistamines (medicines used to treat allergies)
- medicines used to treat epilepsy or fits
- medicines which thin the blood such as warfarin
- other opioid analgesics used to treat pain
- monoamine oxidase inhibitors, medicine used to treat depression, taken within the last 14 days
- flucloxacillin, zidovudine or rifampicin, medicines used to treat infections
- metoclopramide or domperidone, medicines used to control nausea and vomiting
- quinidine, a medicine used to treat abnormal or irregular heartbeat
- phenothiazines and antipsychotic agents, medicines used to treat mental disorders
- medicines used to treat depression
- chloramphenicol, an antibiotic used to treat ear and eye infection
- medicines used to relieve stomach cramps or spasms, to prevent travel sickness and to treat Parkinson's disease
- medicines used to treat high blood pressure
- cholestyramine, a medicine used to lower high cholesterol levels
- chelating resin
- medicines used to treat alcohol and/or opioid dependence (e.g. naltrexone, buprenorphine or methadone)
- medicines for diarrhoea, such as kaolin, pectin and loperamide
- medicines used to control electrolytes levels in kidney disease

These medicines may be affected by Mersyndol DayStrength or may affect how well Mersyndol DayStrength works.

You may need to use different amounts of your medicine, or take different medicines. Your pharmacist or doctor will advise you.

Your doctor or pharmacist will have more information on medicines to be careful with or avoid while taking Mersyndol DayStrength.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Mersyndol DayStrength.

4. How do I use Mersyndol DayStrength?

How much to take

Adults and children 12 years or over:

- One or two tablets every 3 to 4 hours as needed for relief.
- Do not take more than 8 caplets in 24 hour period
- Follow the instructions provided and use Mersyndol DayStrength.

When to take Mersyndol DayStrength

- Mersyndol DayStrength should be used for the temporary relief of acute moderate pain and fever as per the instructions given to you by your doctor.
- Mersyndol DayStrength is not recommended for children under 12 years of age.
- Mersyndol Daystrength is not recommended for use over an extended period of time.
- Do not take more than the recommended dose.

How to take Mersyndol DayStrength

Swallow the tablets whole with a full glass of water or other liquid.

Mersyndol DayStrength can be taken with or without food.

If you forget to use Mersyndol DayStrength

Mersyndol DayStrength should be used exactly as your doctor has instructed.

You may take Mersyndol Daystrength as soon as you remember or if you think you need it.

Do not take a double dose to make up for the dose you have missed.

This may increase the chance of getting unwanted side effect.

If you are not sure what to do, ask your pharmacist or doctor.

If you take too much Mersyndol DayStrength

If you think that you have taken too much Mersyndol DayStrength, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre

(by calling 13 11 26), or

- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

Large amounts of paracetamol, one of the active ingredients, can cause liver damage. Children who take too much Mersyndol DayStrength can also suffer extreme drowsiness, rash or have trouble seeing.

5. What should I know while using Mersyndol DayStrength?

Things you should do

Tell your doctors, dentists and pharmacists who are treating you that you are taking Mersyndol DayStrength.

If you are about to be started on any new medicine, tell your pharmacist or doctor that you are taking Mersyndol DayStrength.

If you plan to have surgery that needs a general anaesthetic, tell your doctor or dentist that you are taking this medicine.

Talk to your pharmacist or doctor about pain control if Mersyndol DayStrength is not helping.

Your pharmacist or doctor will assess your condition and decide if you should continue to take Mersyndol DayStrength.

Things you should not do

Children:

- Do not give this medicine to children under 12 years of age
- Do not give this medicine for more than 48 hours unless a doctor tells you.

Adults:

- Do not take more than a few days at a time unless your doctor tells you to.
- Do not take more than the recommended dose unless your pharmacist or doctor tells you to.
- Do not give this medicine to anyone else, even if they have the same condition as you.
- Do not use this medicine to treat any other complaints unless your pharmacist or doctor tells you to.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Mersyndol DayStrength affects you.

Mersyndol DayStrength may cause drowsiness in some people. If this happens, do not drive or operate machinery.

Drinking alcohol

Tell your doctor if you drink alcohol.

Do not drink while taking Mersyndol DayStrength.

Drinking large quantities of alcohol while taking Mersyndol DayStrength may increase the risk of liver side effects due to paracetamol.

Looking after your medicine

- Store below 30° C.

Follow the instructions in the carton on how to take care of your medicine properly.

Keep your tablets in the blister pack until it's time to take them.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or

- in the car or on windowsills.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

Tell your pharmacist or doctor as soon as possible if you do not feel well while you are taking Mersyndol DayStrength.

Less serious side effects

Less serious side effects	What to do
<p>Stomach and urinary related:</p> <ul style="list-style-type: none"> • constipation • vomiting or nausea • indigestion • difficulty in passing urine <p>Head and neurology related:</p> <ul style="list-style-type: none"> • ringing in the ear • dizziness • drowsiness • dry mouth • headache <p>Allergy related</p> <ul style="list-style-type: none"> • sweating 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p> <p>These are mild side effects of this medicine and usually short-lived.</p>

Serious side effects

Serious side effects	What to do
<p>Allergy related</p> <ul style="list-style-type: none"> • wheezing or difficulty breathing, shallow breathing or shortness of breath, swelling of the face, lips, tongue or other parts of the body • flushing of the face <p>Skin related</p> <ul style="list-style-type: none"> • painful red areas with blisters and peeling layers of skin which may be accompanied by fever and/or chills • severe blisters and bleeding in the lips, eyes, mouth, nose and genitals • skin rashes • Stomach and liver related • hepatitis (symptoms include loss of appetite, itching, yellowing of the skin and eyes, light coloured 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Serious side effects	What to do
bowel motions, dark coloured urine) <ul style="list-style-type: none"> • severe stomach pain, nausea and vomiting Neurological and behavioural <ul style="list-style-type: none"> • unusual or extreme mood swings • dizziness, light-headedness • fast heartbeat 	

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What Mersyndol DayStrength contains

Active ingredient (main ingredient)	Paracetamol and codeine phosphate hemihydrate.
Other ingredients (inactive ingredients)	COMPAP L, sodium starch glycollate, purified talc, magnesium stearate and quinoline yellow (CI 47005).

Do not take this medicine if you are allergic to any of these ingredients.

What Mersyndol DayStrength looks like

Mersyndol DayStrength is available as yellow, capsule-shaped tablets with 'Mersyndol' on one side and 'DS' and a score line on the other (Aust R 59985).

Mersyndol DayStrength is available in packs of 24, 36 and 40 tablets.

Who distributes Mersyndol DayStrength

sanofi-aventis australia Pty Ltd
 12-24 Talavera Road
 Macquarie Park NSW 2113
 Toll Free Number (medical information): 1800 818 806
 Email: medinfo.australia@sanofi.com

This leaflet was prepared in April 2020.

mersyndol-daystrength-ccds4-cmiv11-23apr20